

## **SUPPLEMENTAL MATERIAL**

Genome-wide association trans-ethnic meta-analyses identifies novel associations regulating coagulation Factor VIII and von Willebrand Factor plasma levels

*Maria Sabater-Lleal, PhD; Jennifer E. Huffman, PhD; Paul S. de Vries, PhD; Jonathan Marten et al.*

-Supplementary methods.

-Study descriptions.

-Study specific sources of funding.

-Supplementary tables (note that supplementary Tables S1-S13 are provided in a separate excel file).

Supplementary Table S1: Descriptions of participating cohorts and resources

Supplementary Table S2: Final SNPs composing instrumental variables

Supplementary Table S3: MR results for the instrumental variables individually

Supplementary Table S4: global results for the Egger Regression to evaluate heterogeneity, before and after exclusions

Supplementary Table S5: Results from conditional analyses from European-only data

Supplementary Table S6: Results from conditional analyses from transethnic data

Supplementary Table S7: Haploreg/SNIPA/GWAS catalogue for all transethnic top-SNPs

Supplementary Table S8: eQTL in other tissues

Supplementary Table S9: eQTL in whole blood

Supplementary Table S10: Factor VIII enrichment networks

- Supplementary Table S11: Factor VIII prioritized genes in networks
- Supplementary Table S12: von Willebrand factor enrichment networks
- Supplementary Table S13: von Willebrand factor genes in networks
- Supplementary Figures S1a-n: Regional plots for the novel associations
  - Supplementary Figures S2a-c: Scatter plots for the genetic associations between FVIII and cardiovascular events
  - Supplementary Figure S3: Figure shows the hypothesized effect of the genes found in the present study in relation to the possible regulatory points in VWF synthesis and secretion from endothelial cells. The specific regulatory point suggestion is based on previous literature evidence. Of note, VWF clearance or regulation in platelets were not studied in our in vitro first-pass analyses.

## **SUPPLEMENTARY METHODS**

### *LD-score Regression and Multi-phenotype Analyses*

We assessed the genetic correlation between FVIII and VWF (proportion of shared heritability) using LD-score regression <sup>1</sup> using summary statistics for the European-only meta-analysis of FVIII and VWF (including 25,897 and 42,257 individuals respectively). Given the strong genetic correlation, and in order to gain statistical power, genome-wide autosomal data from the multi-ethnic meta-analysis for FVIII and VWF were combined using aSPU package<sup>2</sup>. Briefly, z-scores for each phenotype were calculated and then used to estimate the covariance between both phenotypes. To overcome the computational burden of running high number of permutations (B) in the whole GWAS dataset, we first run the whole dataset at B=1000, then selected those SNPs where the p-value was smaller than 0.0025 (2.5/number of permutations), and re-run the smaller set of SNPs at increasingly higher B values until the limit of B=10<sup>9</sup> permutations. Since the p-values obtained with this method is

limited by the number of permutations, the final run of  $B=10^9$  permutations limit the minimum p-value to  $2 \times 10^{-9}$ .

#### *Proportion of variance explained*

Proportion of variance  $r^2$  explained by the associated variants was calculated using the formula:

$$r^2 = \frac{2f(1-f)\beta^2}{\text{var}(y)}$$

Where  $f$  is the effect allele frequency and  $\beta$  is the per-allele increment in natural log-transformed phenotype. We calculated beta value using summary-statistics beta values from all participating cohorts and considering the variance,  $\text{var}(y)$ , as square of the weighted mean of standard deviations for the log-transformed phenotype across all cohorts. In addition, we confirmed these values by calculating the proportion of variance explained by the top SNPs in every individual cohort and then summing them up, which resulted in minor differences.

#### *Putative Variants*

Putative functional SNPs for all new loci were investigated *in silico* using publicly available data from Haploreg<sup>3</sup> and SNIIPA<sup>4</sup>.

#### *Expression quantitative trait loci (eQTL) analysis*

The SNP with lowest p-value for each locus was selected and eQTL analysis was performed by checking association with nearby genes (at +/- 1Mb from the selected SNP) in the multi-tissue expression database curated by the Johnson lab at Framingham. A list of all studies included in the database along with their PubMed id is included in below, and results are shown in Supplementary Table 3a and 3b. Results from whole blood and liver studies were prioritized since they have the largest sample size and come from relevant tissues. Since most

studies in the database were not performed using 1000G data, proxy SNPs in high LD ( $r^2 > 0.8$ ) and present in HapMap were selected using SNIPE and also queried.

Database material: A general overview of a subset of >50 eQTL studies has been published<sup>5</sup>, with specific citations for >100 datasets included in the current query following here.

Blood cell related eQTL studies included fresh lymphocytes<sup>6</sup>, fresh leukocytes<sup>7</sup>, leukocyte samples in individuals with Celiac disease<sup>8</sup>, whole blood samples<sup>9-28</sup>, lymphoblastoid cell lines (LCL) derived from asthmatic children<sup>29, 30</sup>, HapMap LCL from 3 populations<sup>31</sup>, a separate study on HapMap CEU LCL<sup>32</sup>, additional LCL population samples<sup>33-39</sup>, neutrophils<sup>40, 41</sup>, CD19+ B cells<sup>42</sup>, primary PHA-stimulated T cells<sup>36, 39</sup>, CD4+ T cells<sup>43</sup>, peripheral blood monocytes<sup>33, 42, 44-47</sup>, long non-coding RNAs in monocytes<sup>48</sup> and CD14+ monocytes before and after stimulation with LPS or interferon-gamma<sup>49</sup>, CD11+ dendritic cells before and after Mycobacterium tuberculosis infection<sup>50</sup> and a separate study of dendritic cells before or after stimulation with LPS, influenza or interferon-beta<sup>51</sup>. Micro-RNA QTLs<sup>52, 53</sup>, DNase-I QTLs<sup>54</sup>, histone acetylation QTLs<sup>55</sup>, and ribosomal occupancy QTLs<sup>56</sup> were also queried for LCL. Splicing QTLs<sup>57</sup> and micro-RNA QTLs<sup>58</sup> were queried in whole blood.

Non-blood cell tissue eQTLs searched included omental and subcutaneous adipose<sup>11, 22, 28, 37, 59</sup>, visceral fat<sup>11</sup>, stomach<sup>59</sup>, endometrial carcinomas<sup>60</sup>, ER+ and ER- breast cancer tumor cells<sup>61</sup>, liver<sup>11, 59, 62-65</sup>, osteoblasts<sup>66</sup>, intestine<sup>67</sup> and normal and cancerous colon<sup>68, 69</sup>, skeletal muscle<sup>11, 70</sup>, breast tissue (normal and cancer)<sup>71, 72</sup>, lung<sup>22, 73-76</sup>, skin<sup>22, 33, 37, 77</sup>, primary fibroblasts<sup>36, 39, 78</sup>, sputum<sup>79</sup>, pancreatic islet cells<sup>80</sup>, prostate<sup>81</sup>, rectal mucosa<sup>82</sup>, arterial wall<sup>11</sup> and heart tissue from left ventricles<sup>22, 83</sup> and left and right atria<sup>84</sup>. Micro-RNA QTLs were

also queried for gluteal and abdominal adipose<sup>85</sup> and liver<sup>86</sup>. Methylation QTLs were queried in pancreatic islet cells<sup>87</sup>. Further mRNA and micro-RNA QTLs were queried from ER+ invasive breast cancer samples, colon-, kidney renal clear-, lung- and prostate-adenocarcinoma samples<sup>88</sup>.

Brain eQTL studies included brain cortex<sup>10, 47, 89-91</sup>, cerebellar cortex<sup>92</sup>, cerebellum<sup>90, 93-96</sup>, frontal cortex<sup>92, 94, 96</sup>, gliomas<sup>97</sup>, hippocampus<sup>92, 94</sup>, inferior olivary nucleus (from medulla), intralobular white matter, and occipital cortex<sup>92</sup>, parietal lobe<sup>95</sup>, pons<sup>96</sup>, pre-frontal cortex<sup>93, 94, 98, 99</sup>, putamen (at the level of anterior commissure)<sup>92</sup>, substantia nigra<sup>92</sup>, temporal cortex<sup>90, 92, 94, 96</sup>, thalamus and visual cortex<sup>94</sup>.

Additional eQTL data was integrated from online sources including ScanDB, the Broad Institute GTEx Portal, and the Pritchard Lab (eqtl.uchicago.edu). Cerebellum, parietal lobe and liver eQTL data was downloaded from ScanDB and cis-eQTLs were limited to those with  $P$ -value  $< 10^{-6}$  and trans-eQTLs with  $P$ -value  $< 5 \times 10^{-8}$ . Results for GTEx Analysis V4 for 13 tissues were downloaded from the GTEx Portal and then additionally filtered as described below [www.gtexportal.org: thyroid, leg skin (sun exposed), tibial nerve, aortic artery, tibial artery, skeletal muscle, esophagus mucosa, esophagus muscularis, lung, heart (left ventricle), stomach, whole blood, and subcutaneous adipose<sup>22</sup>]. Splicing QTL (sQTL) results generated with sQTLseeker with false discovery rate  $P$ -value  $\leq 0.05$  were retained. For all gene-level eQTLs, if at least 1 SNP passed the tissue-specific empirical threshold in GTEx, the best SNP for that eQTL was always retained. All gene-level eQTL SNPs with  $P$ -value  $< 1.67 \times 10^{-11}$  were also retained, reflecting a global threshold correction of  $P$ -value  $= 0.05 / (30,000 \text{ genes} \times 1,000,000 \text{ tests})$ .

### *Pathway analyses*

We used DEPICT (Data-driven Expression-Prioritized Integration for Complex Traits) <sup>100</sup> to infer genes and pathways that were enriched by variants associated with plasma FVIII and VWF. All independent SNPs (defined with  $r^2 < 0.95$  in every 500 Kb region, using 1000G-v3\_GIANT reference data downloaded from <http://csg.sph.umich.edu/abecasis/mach/download/1000G.2012-03-14.html>) with p-value  $< 10^{-5}$  from the European-specific analysis were used as input for the pathway analysis, as recommended in the pipe-line.

### *Mendelian Randomization*

The final variants composing instrumental variables (IV) for FVIII and VWF are listed in Supplementary Table 2a. We conducted MR-Egger regressions <sup>101</sup> and weighted median estimates (WME) <sup>102</sup> to diagnose whether the instrumental variables for the main analysis estimates were valid. To avoid bias due to pleiotropic effect in MR, we visually examined the causal effect estimates produced by each of the individual variants using scatter plots, funnel plots, and forest plots (Supplementary Figure 2), and calculated heterogeneity statistics.<sup>103</sup> This allowed us to select IV as follows: first, we removed the *ABO* and *HLA* loci, which showed known effects with multiple CVD related traits; second, we further removed the *DAB12P* locus for IV of VWF, which was suggested to be an outlier in the heterogeneity test for IVW estimate of VWF effect on CAD; third, we also tested the directional pleiotropic effect using the Egger regression, but no locus was suggested further removed in this step (Supplementary Table 4).

For the multivariate MR approach to test the causal association of FVIII independent of VWF, we used all loci associated with at least one of FVIII and VWF after exclusion strategies. If different sentinel SNPs were reported for FVIII and VWF in the same locus (but

in LD with each other), then a random SNP was selected from these two to represent the locus.

## **STUDY DESCRIPTIONS**

The Age, Gene/Environment Susceptibility-Reykjavik Study (**AGES-Reykjavik**) was initiated in 2002. AGES-Reykjavik was designed to examine risk factors, including genetic susceptibility and gene/environment interaction, in relation to disease and disability in old age <sup>104</sup>. The AGES-Reykjavik sample is drawn from an established population-based cohort, the Reykjavik Study. This cohort of men and women born between 1907 and 1935 has been followed in Iceland since 1967 by the Icelandic Heart Association. The concentration of von Willebrand factor was determined by means of a sensitive enzyme immunoassay <sup>105</sup>.

The **Atherosclerosis Risk in Communities (ARIC)** study has been described in detail previously <sup>106</sup>. Men and women aged 45-64 years at baseline were recruited from four communities: Forsyth County, North Carolina; Jackson, Mississippi; Minneapolis, Minnesota; and Washington County, Maryland. A total of 15,792 individuals, predominantly White and African American, participated in the baseline examination in 1987-1989, with three additional triennial follow-up examinations and a fifth exam in 2011-2013, and a sixth exam in 2016-2017. FVIII activities was measured using a clotting assay (% activity) while VWF antigen was measured using ELISA (% antigen) <sup>107, 108</sup> in plasma samples obtained at the baseline examination.

The **British 1958 birth cohort (B58C)** is a national population sample followed periodically from birth. At age 44-45 years, 9377 cohort members were examined by a

research nurse in the home as described previously and non-fasting blood samples were collected with permission for DNA extraction and creation of immortalised cell cultures<sup>109</sup>. DNA samples from unrelated subjects of white ethnicity, with nationwide geographic coverage, were genotyped either by the Wellcome Trust Case Control Consortium (WTCCC), the Type 1 Diabetes Genetics Consortium, or the GABRIEL consortium<sup>110-112</sup>. Details of the blood collection, VWF measurement and covariate adjustment have been described elsewhere<sup>113</sup>. In brief, VWF antigen was measured by ELISA assays that used a double-antibody sandwich (DAKO, Copenhagen, Denmark). The standard curve was constructed using the 9th British standard for Blood Coagulation Factors from the National Institute for Biological Standards and Controls (NIBSC), South Mimms, Herefordshire UK, and the results were expressed as International units/decilitre (IU/dl). As a control, the pooled plasma of 20 healthy middle-aged persons was run on each ELISA plate. The intra-assay CV was 6%, the inter-assay CV was 8%, and reference range was 50 to 200 IU/dl. Measurements were adjusted for sex, laboratory batch, time of day, month of examination, and postal delay. Adjustment for age was not required as all subjects were aged 44-45 years. Use of anticoagulant therapy was a contraindication to blood sampling. Valid VWF measurements were available for 6093 (93.9%) of the 6491 subjects with genotype data.

**The Coronary Artery Risk Development in Young Adults (CARDIA)** study is a prospective multicenter study with 5115 Caucasian and African American participants ages 18-30 years at baseline, recruited from four centers. The recruitment was done from the total community in Birmingham, AL, from selected census tracts in Chicago, IL and Minneapolis, MN; and from the Kaiser Permanente health plan membership in Oakland, CA. The details of the study design for the CARDIA study have been published before<sup>114</sup>. Nine examinations



have been completed since the baseline examination in 1985–1986, with follow-up examinations 2, 5, 7, 10, 15, 20, 25, and 30 years after baseline. Coagulation FVIII and VWF were measured in plasma citrate <sup>115</sup>. FVIII coagulant activities were assayed by a one-stage system with reagents from Pacific Hemostasis and George King Biomedical, Inc. The standard curve was prepared by using universal reference plasma from Curtin Matheson Scientific and the results calculated as a percentage of standard with the data management system of the MLA Electra 800. von Willebrand antigen was measured by an enzyme-linked immunosorbent assay obtained from American Bioproducts Co <sup>116</sup>.

The **Cardiovascular Health Study (CHS)** is a population-based cohort study of risk factors for coronary heart disease and stroke in adults  $\geq 65$  years conducted across four field centers <sup>117</sup>. The original predominantly European -ancestry cohort of 5,201 persons was recruited in 1989-1990 from random samples of the Medicare eligibility lists; subsequently, an additional predominantly African-American cohort of 687 persons were enrolled for a total sample of 5,888. Blood samples were drawn from all participants at their baseline examination and DNA was subsequently extracted from available samples. Genotyping was performed at the General Clinical Research Center's Phenotyping/Genotyping Laboratory at Cedars-Sinai among CHS participants who consented to genetic testing and had DNA available using the Illumina 370CNV BeadChip system (for European ancestry participants, in 2007) or the Illumina HumanOmni1-Quad\_v1 BeadChip system (for African-American participants, in 2010). CHS was approved by institutional review committees at each field center.

Participants included in the present analyses had available DNA and gave informed consent including consent to use of genetic information for the study of cardiovascular disease. FVIII activity determined by using factor VIII-deficient plasma (Organon-Teknika) and partial thromboplastin (Organon-Teknika) <sup>118</sup>. Unassayed pooled normal plasma (George King

Biomedical, Overland Park, KS, USA) was used as the standard and calibrated with the World Health Organization reference plasma for both assays. Activity measured using Coag-A-Mate X2 (Organon-Teknika). Values were expressed as percentage of the standard. The intra-assay CV was 12.6% and inter-assay CV was 13.8%. No reference range available.

The **CROATIA-Vis** study, Croatia, is a family-based, cross-sectional study in the isolated island of Vis that included 1,056 examinees aged 18-93. Blood samples were collected in 2003 and 2004 along with many clinical and biochemical measures and lifestyle and health questionnaires. The VWF antigen was measured by ELISA assays that used a double-antibody sandwich (DAKO, Copenhagen, Denmark).

The **Framingham Heart Study (FHS)** was started in 1948 with 5,209 randomly ascertained participants from Framingham, Massachusetts, US, who had undergone biannual examinations to investigate cardiovascular disease and its risk factors. In 1971, the Offspring cohort (comprising 5,124 children of the original cohort and the children's spouses) and in 2002, the Third Generation (consisting of 4,095 children of the Offspring cohort) were recruited. FHS participants in this study are of European ancestry. The methods of recruitment and data collection for the Offspring and Third Generation cohorts have been described elsewhere <sup>119</sup>. Von Willebrand factor was assessed using ELISA at exam 5 (1991-1995) in the Offspring cohort. In our laboratory, the intra-assay coefficient of variation was 8.8%. No reference range is available <sup>120</sup>.

The **Genes and Blood-Clotting Study (GABC)** consists of a cohort of 1,150 healthy siblings recruited from the University of Michigan, Ann Arbor, between June 26, 2006 and January 30, 2009. Participants were between the ages of 14 and 35 y, and had at least one eligible

healthy sibling. Subjects who indicated that they were pregnant, had a known bleeding or blood-clotting disorder, or any illness requiring regular medical care were excluded. All participants provided informed consent. Subjects completed an online phenotyping survey and donated a blood sample for DNA extraction and plasma biochemical phenotyping. VWF levels were determined at the University of Michigan using a custom AlphaLISA™ assay (Perkin-Elmer), which utilized a polyclonal anti-von Willebrand factor antibody from DAKO Cytomation.

The **GAIT (Genetic Analysis of Idiopathic Thrombophilia)** project is a family based study where 935 subjects in 35 extended pedigrees were collected. To be included in the study, a family was required to have at least 10 living individuals in 3 or more generations. Families were selected through a proband with idiopathic thrombophilia, which was defined as recurrent thrombotic events (at least one of which was spontaneous), a single spontaneous thrombotic episode plus a first-degree relative also affected, or onset of thrombosis before age 45. Thrombosis in these probands was considered idiopathic when biological causes as antithrombin deficiency, protein S and C deficiencies, activated protein C resistance, plasminogen deficiency, heparin cofactor II deficiency, Factor V Leiden, dysfibrinogenemia, lupus anticoagulant and antiphospholipid antibodies, were excluded. Subjects were interviewed by a physician to determine their health and reproductive history, current medications, alcohol consumption, use of sex hormones (oral contraceptives or hormonal replacement therapy) and their smoking history. The study was performed according to the Declaration of Helsinki. All procedures of the study were reviewed by the Institutional Review Board of the Hospital de la Santa Creu i Sant Pau, Barcelona, Spain. Adult subjects gave informed consent for themselves and for their minor children. Coagulation factors were measured in platelet-poor citrated plasma<sup>121</sup>. Coagulation FVIII was assayed with deficient

plasma from Diagnostica Stago (Asnieres). von Willebrand factor was measured by an ELISA method with antibodies from Dako. To reduce measurement error, assays were performed in duplicate, and the average value was calculated for each person. Intra-assay and interassay coefficients of variation were generally estimated to be between 2% and 6%.

**GeneSTAR (Genetic Study of Atherosclerosis Risk)** is an ongoing prospective family study begun in 1983 to explore the causes of early-onset cardiovascular disease. Probands with an early-onset coronary disease event prior to 60 years of age were identified at the time of hospitalization in any of 10 Baltimore area hospitals. Their apparently healthy 30-59 year old siblings without known CAD were initially recruited and screened between 1983 and 2006; offspring of the siblings and probands, as well as the co-parent of these offspring, were recruited and assessed between 2003 and 2006. In this study, European and African American participants with both genotyping with the Illumina Human1Mv1\_C chip and measured VWF were included. Levels of VWF were measured using ELISA at the University of Maryland Cytokine Laboratory.

**The Ludwigshafen Risk and Cardiovascular Health (LURIC)** study is a monocentric hospital based prospective study including 3316 individuals referred for coronary angiography recruited in the Ludwigshafen Cardiac Center, southwestern Germany from 1997 – 2000. Clinical indications for angiography were chest pain or a positive non-invasive stress test suggestive of myocardial ischemia. To limit clinical heterogeneity, individuals suffering from acute illnesses other than acute coronary syndrome, chronic non-cardiac diseases and a history of malignancy within the five past years were excluded. All participants were of European ancestry and completed a detailed questionnaire which gathered information on medical history, clinical, and lifestyle factors. Study protocols were

approved by the ethics committee of the "Landesärztekammer Rheinland-Pfalz" and the study was conducted in accordance with the "Declaration of Helsinki". Informed written consent was obtained from all participants. Coronary heart disease at baseline was defined as the presence of a visible luminal narrowing ( $>50\%$  stenosis) in at least one of 15 coronary segments according to the classification of the American Heart Association. Fasting blood samples were obtained by venipuncture in the early morning and stored for later analyses. Coagulation FVIII and VWF were measured in plasma using methods previously described<sup>122</sup>. For this study a subset of 3061 Samples were used that had been genotyped on an Affymetrix 6.0 array.

The **MARseille THrombosis Association (MARTHA)** project has already been extensively<sup>123</sup>. It is composed of unrelated subjects of European origin, with the majority being of French ancestry, consecutively recruited at the Thrombophilia center of La Timone hospital (Marseille, France) between January 1994 and October 2012. All patients had a documented history of VT and free of well characterized genetic risk factors including AT, PC, or PS deficiency, homozygosity for FV Leiden or FII 20210A, and lupus anticoagulant. They were interviewed by a physician on their medical history, which emphasized manifestations of deep vein thrombosis and pulmonary embolism using a standardized questionnaire. The thrombotic events were confirmed by venography, Doppler ultrasound, spiral computed tomographic scanning angiography, and/or ventilation/perfusion lung scan. Hemostasis-related parameters were centrally performed using the Star automate and commercially available kits and reagents from Diagnostica Stago (Asnières, France) including the corresponding normal and pathological control plasmas and standard plasmas. FVIII and VWF levels are respectively measured by VIII:C using human FVIII-deficient plasma in a 1-stage factor assay and by STA LIATEST VWF (Diagnostica Stago) on Star automate. They

were measured at least 3 months after the most recent VT event to minimize the effect of the acute phase <sup>124</sup>. From the 1542 MARTHA participants with GWAS data, 727 and 877 were included for the present analyses on FVIII and VWF levels.

**The Multiple Environmental and Genetic Assessment of risk factors for venous thrombosis (MEGA)** study is a large population-based case-control study <sup>125</sup>. Data collection and ascertainment of venous thrombotic events have been previously described in detail. In short, patients with a first deep vein thrombosis or pulmonary embolism were recruited at six anticoagulation clinics in the Netherlands between 1999 and 2004. The diagnosis of a deep vein thrombosis was based on compression ultrasonography, whereas a pulmonary embolism was confirmed by perfusion and ventilation scintigraphy, helical computed tomography or pulmonary angiography. Blood samples were taken at least 3 months after discontinuation of vitamin K antagonist treatment, unless patients were still receiving anticoagulant therapy one year after their VT event. For the present analyses, patients who were still receiving anticoagulant treatment at the time of blood collection were excluded. Factor VIII activity levels were measured with a mechanical clot detection method on an STA-R coagulation analyzer (Diagnostica Stago, Asnieres, France), whereas von Willebrand factor antigen levels were measured with the immunoturbidimetric method using the STA liatest kit (rabbit anti-human von Willebrand factor antibodies). For genome-wide genotyping with the Illumina Human660-Quad Beadchip, we sampled 1,499 patients with a first episode of VT. Patients with a cancer diagnosis were excluded. Patients with genotyping success lower than 95% were excluded from the analyses as were patients showing discrepancies between their reported and genotypic sex. Individuals demonstrating a too high or low level of heterozygosity, close relatedness or non-European ancestry were also excluded. Variants showing significant ( $P\text{-value} < 1 \times 10^{-6}$ )

deviation from HWE, with MAF less than 1%, and with genotyping call rate <98% were filtered out. After quality-control, Imputation to the 1000 Genomes population reference (Phase1 March 2012 release) was performed using IMPUTE2 software. Association analyses were conducted with the score test as implemented in SNPTEST, adjusting for age, sex, and, the first three principal components.

**The Multi-Ethnic Study of Atherosclerosis (MESA)** is a study of the characteristics of subclinical cardiovascular disease and the risk factors that predict progression to clinically overt cardiovascular disease or progression of the subclinical disease<sup>126</sup>. MESA consisted of a diverse, population-based sample of an initial 6,814 asymptomatic men and women aged 45-84. 38 percent of the recruited participants were white, 28 percent African American, 22 percent Hispanic, and 12 percent Asian, predominantly of Chinese descent. Participants were recruited from six field centers across the United States: Wake Forest University, Columbia University, Johns Hopkins University, University of Minnesota, Northwestern University and University of California - Los Angeles. The first examination took place over two years, from July 2000 - July 2002. Only individuals without clinical CVD were eligible. Each participant received an extensive physical exam and determination of coronary calcification, ventricular mass and function, flow-mediated endothelial vasodilation, carotid intimal-medial wall thickness and presence of echogenic lucencies in the carotid artery, lower extremity vascular insufficiency, arterial wave forms, electrocardiographic (ECG) measures, standard coronary risk factors, sociodemographic factors, lifestyle factors, and psychosocial factors. It was followed by four examination periods that were 17-20 months in length, and a sixth exam is currently taking place. Factor VIII levels were determined by measuring the clot time of a sample in factor VIII deficient plasma in the presence of activators utilizing the Sta-R analyzer (STA-Deficient VIII; Diagnostica Stago, Parsippany, NJ). The assay was performed

at the Laboratory for Clinical Biochemistry Research (University of Vermont, Burlington, VT). Results are given as percent factor VIII. The normal plasma range of factor VIII in healthy adults ranges from 60-150%. The analytical coefficient of variation for the factor VIII assay was 10%. von Willebrand factor (VWF) was measured by immunoturbidimetric assay on the Sta-R analyzer (Iatest VWF; Diagnostica Stago, Parsippany, NJ). The assay utilizes latex particles to which specific antibodies have been attached. In the presence of antigen (VWF), the particles agglutinate to form aggregates, which absorb more light. This increase in absorbance is proportional to the VWF present in the test sample. This assay was performed at the Laboratory for Clinical Biochemistry Research (University of Vermont, Burlington, VT). The results are presented as percent VWF. The expected range in healthy adults is 50-160%. Intra- and inter-assay analytical coefficients of variation were 3.7% and 4.5%, respectively.

The **Orkney Complex Disease Study (ORCADES)** is a family-based, cross-sectional study in the isolated Scottish archipelago of Orkney. Genetic diversity in this population is decreased compared to Mainland Scotland, consistent with the high levels of endogamy historically. Data for participants aged 18-100 years, from a subgroup of ten islands, were used for this analysis. Fasting blood samples were collected and over 300 health-related phenotypes and environmental exposures were measured in each individual. All participants gave informed consent and the study was approved by Research Ethics Committees in Orkney and Aberdeen. Subjects with self-reported non-European ancestry were excluded. The VWF antigen was measured by ELISA assays that used a double-antibody sandwich (DAKO, Copenhagen, Denmark).

The **PROspective Study of Pravastatin in the Elderly at Risk (PROSPER)** was a



prospective multicenter randomized placebo-controlled trial to assess whether treatment with pravastatin diminishes the risk of major vascular events in elderly. Between December 1997 and May 1999, subjects were enrolled in Scotland (Glasgow), Ireland (Cork), and the Netherlands (Leiden). Men and women aged 70-82 years were recruited if they had pre-existing vascular disease or increased risk of such disease because of smoking, hypertension, or diabetes. A total number of 5,804 subjects were randomly assigned to pravastatin or placebo. A large number of prospective tests were performed including Biobank tests and cognitive function measurements. A whole genome wide screening has been performed in the sequential PHASE project with the use of the Illumina 660K beadchip. Of 5,763 subjects DNA was available for genotyping. Genotyping was performed with the Illumina 660K beadchip, after QC (call rate <95%) 5,244 subjects and 557,192 SNPs were left for analysis. Imputation was based on the March 2012 1000G release using the IMPUTE2 imputation software. For this study, 4927 PROSPER participants with data on VWF levels in citrate (ELISA) were analyzed.

The **Rotterdam Study (RS-I and RS-II)** is a prospective, population-based cohort study of determinants of several chronic diseases in older adults <sup>127</sup>. RS-I comprised 7,983 inhabitants of Ommoord, a district of Rotterdam in the Netherlands, who were 55 years or over. The baseline examination took place between 1990 and 1993. In 1999, the cohort was extended to include 3011 inhabitants who reached the age of 55 years after the baseline examination and persons aged 55 years or older who migrated into the research area (RS-II). Subjects are of European ancestry based on their self-report. Factor VIII activity was measured with a one-stage clotting assay by using a mixture of micronized silica and phospholipids (Platelin LS, Biomerieux) and factor VIII-deficient plasma (Biopool). The plasma concentrations were expressed as percentage activity by relating the clotting time to a calibration curve

constructed of a standardized control plasma. As a control, the pooled plasma of 50 healthy middle-aged persons was used and three control samples were run with each batch of study samples. The intra-assay CV was 2.9%, the inter-assay CV was 5.2%, and the reference range was 0.70-1.40 U/ml. Von Willebrand factor antigen was measured with an in-house ELISA with polyclonal rabbit anti-human VWF antibodies (DAKO). The intra-assay CV was 1.9%, inter-assay CV was 6.3%, and the reference range was 0.60-1.40 U/ml.

The **Trinity Student Study (TSS)** consists of a cohort of 2,524 healthy, ethnically Irish individuals, attending the University of Dublin, Trinity College, with ages between 18 and 28 years, recruited over one academic year in 2003–2004. Ethical approval was obtained from the Dublin Federated Hospitals Research Ethics Committee, which is affiliated with the Trinity College, and reviewed by the Office of Human Subjects Research at the National Institutes of Health. Written informed consent was obtained from participants before recruitment. VWF levels were determined at the University of Michigan using a custom AlphaLISA™ assay (Perkin-Elmer), which utilized a polyclonal anti-von Willebrand factor antibody from DAKO Cytomation.

## REFERENCES:

1. Bulik-Sullivan BK, Loh PR, Finucane HK, Ripke S, Yang J, Schizophrenia Working Group of the Psychiatric Genomics Consortium, Patterson N, Daly MJ, Price AL and Neale BM. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet.* 2015;47:291-5.
2. Kwak IY and Pan W. Adaptive gene- and pathway-trait association testing with GWAS summary statistics. *Bioinformatics.* 2016;32:1178-84.
3. Ward LD and Kellis M. HaploReg v4: systematic mining of putative causal variants, cell types, regulators and target genes for human complex traits and disease. *Nucleic Acids Res.* 2016;44:D877-81.
4. Arnold M, Raffler J, Pfeufer A, Suhre K and Kastenmuller G. SNIIPA: an interactive, genetic variant-centered annotation browser. *Bioinformatics.* 2015;31:1334-6.
5. Zhang X, Gierman HJ, Levy D, Plump A, Dobrin R, Goring HH, Curran JE, Johnson MP, Blangero J, Kim SK, O'Donnell CJ, Emilsson V and Johnson AD. Synthesis of 53 tissue

and cell line expression QTL datasets reveals master eQTLs. *BMC Genomics*. 2014;15:532.

6. Goring HH, Curran JE, Johnson MP, Dyer TD, Charlesworth J, Cole SA, Jowett JB, Abraham LJ, Rainwater DL, Comuzzie AG, Mahaney MC, Almasy L, MacCluer JW, Kissebah AH, Collier GR, Moses EK and Blangero J. Discovery of expression QTLs using large-scale transcriptional profiling in human lymphocytes. *Nat Genet*. 2007;39:1208-16.
7. Idaghdour Y, Czika W, Shianna KV, Lee SH, Visscher PM, Martin HC, Miclaus K, Jadallah SJ, Goldstein DB, Wolfinger RD and Gibson G. Geographical genomics of human leukocyte gene expression variation in southern Morocco. *Nat Genet*. 2010;42:62-7.
8. Heap GA, Trynka G, Jansen RC, Bruinenberg M, Swertz MA, Dinesen LC, Hunt KA, Wijmenga C, Vanheel DA and Franke L. Complex nature of SNP genotype effects on gene expression in primary human leucocytes. *BMC Med Genomics*. 2009;2:1.
9. Chen W, Brehm JM, Lin J, Wang T, Forno E, Acosta-Perez E, Boutaoui N, Canino G and Celedon JC. Expression quantitative trait loci (eQTL) mapping in Puerto Rican children. *PLoS One*. 2015;10:e0122464.
10. Lock EF, Soldano KL, Garrett ME, Cope H, Markunas CA, Fuchs H, Grant G, Dunson DB, Gregory SG and Ashley-Koch AE. Joint eQTL assessment of whole blood and dura mater tissue from individuals with Chiari type I malformation. *BMC Genomics*. 2015;16:11.
11. Foroughi Asl H, Talukdar HA, Kindt AS, Jain RK, Ermel R, Ruusalepp A, Nguyen KD, Dobrin R, Reilly DF, Schunkert H, Samani NJ, Braenne I, Erdmann J, Melander O, Qi J, Ivert T, Skogsberg J, Schadt EE, Michoel T, Bjorkegren JL and Consortium CA. Expression quantitative trait Loci acting across multiple tissues are enriched in inherited risk for coronary artery disease. *Circ Cardiovasc Genet*. 2015;8:305-15.
12. Pierce BL, Tong L, Chen LS, Rahaman R, Argos M, Jasmine F, Roy S, Paul-Brutus R, Westra HJ, Franke L, Esko T, Zaman R, Islam T, Rahman M, Baron JA, Kibriya MG and Ahsan H. Mediation analysis demonstrates that trans-eQTLs are often explained by cis-mediation: a genome-wide analysis among 1,800 South Asians. *PLoS Genet*. 2014;10:e1004818.
13. Narahara M, Higasa K, Nakamura S, Tabara Y, Kawaguchi T, Ishii M, Matsubara K, Matsuda F and Yamada R. Large-scale East-Asian eQTL mapping reveals novel candidate genes for LD mapping and the genomic landscape of transcriptional effects of sequence variants. *PLoS One*. 2014;9:e100924.
14. Schramm K, Marzi C, Schurmann C, Carstensen M, Reinmaa E, Biffar R, Eckstein G, Gieger C, Grabe HJ, Homuth G, Kastenmuller G, Magi R, Metspalu A, Mihailov E, Peters A, Petersmann A, Roden M, Strauch K, Suhre K, Teumer A, Volker U, Volzke H, Wang-Sattler R, Waldenberger M, Meitinger T, Illig T, Herder C, Grallert H and Prokisch H. Mapping the genetic architecture of gene regulation in whole blood. *PLoS One*. 2014;9:e93844.
15. Wright FA, Sullivan PF, Brooks AI, Zou F, Sun W, Xia K, Madar V, Jansen R, Chung W, Zhou YH, Abdellaoui A, Batista S, Butler C, Chen G, Chen TH, D'Ambrosio D, Gallins P, Ha MJ, Hottenga JJ, Huang S, Kattenberg M, Kochar J, Middeldorp CM, Qu A, Shabalin A, Tischfield J, Todd L, Tzeng JY, van Grootheest G, Vink JM, Wang Q, Wang W, Wang W, Willemsen G, Smit JH, de Geus EJ, Yin Z, Penninx BW and Boomsma DI. Heritability and genomics of gene expression in peripheral blood. *Nat Genet*. 2014;46:430-7.
16. Quinlan J, Idaghdour Y, Goulet JP, Gbeha E, de Malliard T, Bruat V, Grenier JC, Gomez S, Sanni A, Rahimy MC and Awadalla P. Genomic architecture of sickle cell disease in West African children. *Front Genet*. 2014;5:26.

17. Benton MC, Lea RA, Macartney-Coxson D, Carless MA, Goring HH, Bellis C, Hanna M, Eccles D, Chambers GK, Curran JE, Harper JL, Blangero J and Griffiths LR. Mapping eQTLs in the Norfolk Island genetic isolate identifies candidate genes for CVD risk traits. *Am J Hum Genet.* 2013;93:1087-99.
18. Battle A, Mostafavi S, Zhu X, Potash JB, Weissman MM, McCormick C, Haudenschild CD, Beckman KB, Shi J, Mei R, Urban AE, Montgomery SB, Levinson DF and Koller D. Characterizing the genetic basis of transcriptome diversity through RNA-sequencing of 922 individuals. *Genome Res.* 2014;24:14-24.
19. Westra HJ, Peters MJ, Esko T, Yaghootkar H, Schurmann C, Kettunen J, Christiansen MW, Fairfax BP, Schramm K, Powell JE, Zhernakova A, Zhernakova DV, Veldink JH, Van den Berg LH, Karjalainen J, Withoff S, Uitterlinden AG, Hofman A, Rivadeneira F, Hoen PAC, Reinmaa E, Fischer K, Nelis M, Milani L, Melzer D, Ferrucci L, Singleton AB, Hernandez DG, Nalls MA, Homuth G, Nauck M, Radke D, Volker U, Perola M, Salomaa V, Brody J, Suchy-Dicey A, Gharib SA, Enquobahrie DA, Lumley T, Montgomery GW, Makino S, Prokisch H, Herder C, Roden M, Grallert H, Meitinger T, Strauch K, Li Y, Jansen RC, Visscher PM, Knight JC, Psaty BM, Ripatti S, Teumer A, Frayling TM, Metspalu A, van Meurs JBJ and Franke L. Systematic identification of trans eQTLs as putative drivers of known disease associations. *Nat Genet.* 2013;45:1238-1243.
20. Landmark-Hoyvik H, Dumeaux V, Nebdal D, Lund E, Tost J, Kamatani Y, Renault V, Borresen-Dale AL, Kristensen V and Edvardsen H. Genome-wide association study in breast cancer survivors reveals SNPs associated with gene expression of genes belonging to MHC class I and II. *Genomics.* 2013;102:278-87.
21. Zhernakova DV, de Klerk E, Westra HJ, Mastrokolas A, Amini S, Ariyurek Y, Jansen R, Penninx BW, Hottenga JJ, Willemsen G, de Geus EJ, Boomsma DI, Veldink JH, van den Berg LH, Wijmenga C, den Dunnen JT, van Ommen GJ, t Hoen PA and Franke L. DeepSAGE reveals genetic variants associated with alternative polyadenylation and expression of coding and non-coding transcripts. *PLoS Genet.* 2013;9:e1003594.
22. Consortium GT. The Genotype-Tissue Expression (GTEx) project. *Nat Genet.* 2013;45:580-5.
23. Sasayama D, Hori H, Nakamura S, Miyata R, Teraishi T, Hattori K, Ota M, Yamamoto N, Higuchi T, Amano N and Kunugi H. Identification of single nucleotide polymorphisms regulating peripheral blood mRNA expression with genome-wide significance: an eQTL study in the Japanese population. *PLoS One.* 2013;8:e54967.
24. van Eijk KR, de Jong S, Boks MP, Langeveld T, Colas F, Veldink JH, de Kovel CG, Janson E, Strengman E, Langfelder P, Kahn RS, van den Berg LH, Horvath S and Ophoff RA. Genetic analysis of DNA methylation and gene expression levels in whole blood of healthy human subjects. *BMC Genomics.* 2012;13:636.
25. Mehta D, Heim K, Herder C, Carstensen M, Eckstein G, Schurmann C, Homuth G, Nauck M, Volker U, Roden M, Illig T, Gieger C, Meitinger T and Prokisch H. Impact of common regulatory single-nucleotide variants on gene expression profiles in whole blood. *Eur J Hum Genet.* 2013;21:48-54.
26. Powell JE, Henders AK, McRae AF, Caracella A, Smith S, Wright MJ, Whitfield JB, Dermitzakis ET, Martin NG, Visscher PM and Montgomery GW. The Brisbane Systems Genetics Study: genetical genomics meets complex trait genetics. *PLoS One.* 2012;7:e35430.
27. Fehrmann RS, Jansen RC, Veldink JH, Westra HJ, Arends D, Bonder MJ, Fu J, Deelen P, Groen HJ, Smolonska A, Weersma RK, Hofstra RM, Buurman WA, Rensen S, Wolfs MG, Platteel M, Zhernakova A, Elbers CC, Festen EM, Trynka G, Hofker MH, Saris

- CG, Ophoff RA, van den Berg LH, van Heel DA, Wijmenga C, Te Meerman GJ and Franke L. Trans-eQTLs reveal that independent genetic variants associated with a complex phenotype converge on intermediate genes, with a major role for the HLA. *PLoS Genet.* 2011;7:e1002197.
28. Emilsson V, Thorleifsson G, Zhang B, Leonardson AS, Zink F, Zhu J, Carlson S, Helgason A, Walters GB, Gunnarsdottir S, Mouy M, Steinthorsdottir V, Eiriksdottir GH, Bjornsdottir G, Reynisdottir I, Gudbjartsson D, Helgadottir A, Jonasdottir A, Jonasdottir A, Styrkarsdottir U, Gretarsdottir S, Magnusson KP, Stefansson H, Fossdal R, Kristjansson K, Gislason HG, Stefansson T, Leifsson BG, Thorsteinsdottir U, Lamb JR, Gulcher JR, Reitman ML, Kong A, Schadt EE and Stefansson K. Genetics of gene expression and its effect on disease. *Nature.* 2008;452:423-8.
29. Liang L, Morar N, Dixon AL, Lathrop GM, Abecasis GR, Moffatt MF and Cookson WO. A cross-platform analysis of 14,177 expression quantitative trait loci derived from lymphoblastoid cell lines. *Genome Res.* 2013;23:716-26.
30. Dixon AL, Liang L, Moffatt MF, Chen W, Heath S, Wong KC, Taylor J, Burnett E, Gut I, Farrall M, Lathrop GM, Abecasis GR and Cookson WO. A genome-wide association study of global gene expression. *Nat Genet.* 2007;39:1202-7.
31. Stranger BE, Nica AC, Forrest MS, Dimas A, Bird CP, Beazley C, Ingle CE, Dunning M, Flicek P, Koller D, Montgomery S, Tavare S, Deloukas P and Dermitzakis ET. Population genomics of human gene expression. *Nat Genet.* 2007;39:1217-24.
32. Kwan T, Benovoy D, Dias C, Gurd S, Provencher C, Beaulieu P, Hudson TJ, Sladek R and Majewski J. Genome-wide analysis of transcript isoform variation in humans. *Nat Genet.* 2008;40:225-31.
33. Huang J, Chen J, Esparza J, Ding J, Elder JT, Abecasis GR, Lee YA, Mark Lathrop G, Moffatt MF, Cookson WO and Liang L. eQTL mapping identifies insertion- and deletion-specific eQTLs in multiple tissues. *Nature communications.* 2015;6:6821.
34. Bryois J, Buil A, Evans DM, Kemp JP, Montgomery SB, Conrad DF, Ho KM, Ring S, Hurles M, Deloukas P, Davey Smith G and Dermitzakis ET. Cis and trans effects of human genomic variants on gene expression. *PLoS Genet.* 2014;10:e1004461.
35. Mangravite LM, Engelhardt BE, Medina MW, Smith JD, Brown CD, Chasman DI, Mecham BH, Howie B, Shim H, Naidoo D, Feng Q, Rieder MJ, Chen YD, Rotter JL, Ridker PM, Hopewell JC, Parish S, Armitage J, Collins R, Wilke RA, Nickerson DA, Stephens M and Krauss RM. A statin-dependent QTL for GATM expression is associated with statin-induced myopathy. *Nature.* 2013;502:377-80.
36. Gutierrez-Arcelus M, Lappalainen T, Montgomery SB, Buil A, Ongen H, Yurovsky A, Bryois J, Giger T, Romano L, Planchon A, Falconnet E, Bielser D, Gagnebin M, Padioulet I, Borel C, Letourneau A, Makrythanasis P, Guipponi M, Gehrig C, Antonarakis SE and Dermitzakis ET. Passive and active DNA methylation and the interplay with genetic variation in gene regulation. *Elife.* 2013;2:e00523.
37. Grundberg E, Small KS, Hedman AK, Nica AC, Buil A, Keildson S, Bell JT, Yang TP, Meduri E, Barrett A, Nisbett J, Sekowska M, Wilk A, Shin SY, Glass D, Travers M, Min JL, Ring S, Ho K, Thorleifsson G, Kong A, Thorsteindottir U, Ainali C, Dimas AS, Hassanali N, Ingle C, Knowles D, Krestyaninova M, Lowe CE, Di Meglio P, Montgomery SB, Parts L, Potter S, Surdulescu G, Tsaprouni L, Tsoka S, Bataille V, Durbin R, Nestle FO, O'Rahilly S, Soranzo N, Lindgren CM, Zondervan KT, Ahmadi KR, Schadt EE, Stefansson K, Smith GD, McCarthy MI, Deloukas P, Dermitzakis ET, Spector TD and Multiple Tissue Human Expression Resource C. Mapping cis- and trans-regulatory effects across multiple tissues in twins. *Nat Genet.* 2012;44:1084-9.

38. Cusanovich DA, Billstrand C, Zhou X, Chavarria C, De Leon S, Michelini K, Pai AA, Ober C and Gilad Y. The combination of a genome-wide association study of lymphocyte count and analysis of gene expression data reveals novel asthma candidate genes. *Hum Mol Genet.* 2012;21:2111-23.
39. Dimas AS, Deutsch S, Stranger BE, Montgomery SB, Borel C, Attar-Cohen H, Ingle C, Beazley C, Gutierrez Arcelus M, Sekowska M, Gagnebin M, Nisbett J, Deloukas P, Dermitzakis ET and Antonarakis SE. Common regulatory variation impacts gene expression in a cell type-dependent manner. *Science.* 2009;325:1246-50.
40. Andiappan AK, Melchioni R, Poh TY, Nah M, Puan KJ, Vigano E, Haase D, Yusof N, San Luis B, Lum J, Kumar D, Foo S, Zhuang L, Vasudev A, Irwanto A, Lee B, Nardin A, Liu H, Zhang F, Connolly J, Liu J, Mortellaro A, Wang de Y, Poidinger M, Larbi A, Zolezzi F and Rotzschke O. Genome-wide analysis of the genetic regulation of gene expression in human neutrophils. *Nature communications.* 2015;6:7971.
41. Naranbhai V, Fairfax BP, Makino S, Humburg P, Wong D, Ng E, Hill AV and Knight JC. Genomic modulators of gene expression in human neutrophils. *Nature communications.* 2015;6:7545.
42. Fairfax BP, Makino S, Radhakrishnan J, Plant K, Leslie S, Dilthey A, Ellis P, Langford C, Vannberg FO and Knight JC. Genetics of gene expression in primary immune cells identifies cell type-specific master regulators and roles of HLA alleles. *Nat Genet.* 2012;44:502-10.
43. Murphy A, Chu JH, Xu M, Carey VJ, Lazarus R, Liu A, Szeffler SJ, Strunk R, Demuth K, Castro M, Hansel NN, Diette GB, Vonakis BM, Adkinson NF, Jr., Klanderman BJ, Senter-Sylvia J, Ziniti J, Lange C, Pastinen T and Raby BA. Mapping of numerous disease-associated expression polymorphisms in primary peripheral blood CD4+ lymphocytes. *Hum Mol Genet.* 2010;19:4745-57.
44. Kirsten H, Al-Hasani H, Holdt L, Gross A, Beutner F, Krohn K, Horn K, Ahnert P, Burkhardt R, Reiche K, Hackermuller J, Loffler M, Teupser D, Thiery J and Scholz M. Dissecting the genetics of the human transcriptome identifies novel trait-related trans-eQTLs and corroborates the regulatory relevance of non-protein coding locidagger. *Hum Mol Genet.* 2015;24:4746-63.
45. Almlöf JC, Lundmark P, Lundmark A, Ge B, Maouche S, Goring HH, Liljedahl U, Enström C, Brocheton J, Proust C, Godefroy T, Sambrook JG, Jolley J, Crisp-Hihn A, Foad N, Lloyd-Jones H, Stephens J, Gwilliam R, Rice CM, Hengstenberg C, Samani NJ, Erdmann J, Schunkert H, Pastinen T, Deloukas P, Goodall AH, Ouwehand WH, Cambien F and Syvänen AC. Powerful identification of cis-regulatory SNPs in human primary monocytes using allele-specific gene expression. *PLoS One.* 2012;7:e52260.
46. Zeller T, Wild P, Szymczak S, Rotival M, Schillert A, Castagne R, Maouche S, Germain M, Lackner K, Rossmann H, Eleftheriadis M, Sinning CR, Schnabel RB, Lubos E, Mennerich D, Rust W, Perret C, Proust C, Nicaud V, Loscalzo J, Hubner N, Tregouet D, Munzel T, Ziegler A, Tiret L, Blankenberg S and Cambien F. Genetics and beyond--the transcriptome of human monocytes and disease susceptibility. *PLoS One.* 2010;5:e10693.
47. Heinzen EL, Ge D, Cronin KD, Maia JM, Shianna KV, Gabriel WN, Welsh-Bohmer KA, Hulette CM, Denny TN and Goldstein DB. Tissue-specific genetic control of splicing: implications for the study of complex traits. *PLoS Biol.* 2008;6:e1.
48. Almlöf JC, Lundmark P, Lundmark A, Ge B, Pastinen T, Cardiogenics C, Goodall AH, Cambien F, Deloukas P, Ouwehand WH and Syvänen AC. Single nucleotide polymorphisms with cis-regulatory effects on long non-coding transcripts in human primary monocytes. *PLoS One.* 2014;9:e102612.

49. Fairfax BP, Humburg P, Makino S, Naranbhai V, Wong D, Lau E, Jostins L, Plant K, Andrews R, McGee C and Knight JC. Innate immune activity conditions the effect of regulatory variants upon monocyte gene expression. *Science*. 2014;343:1246949.
50. Barreiro LB, Tailleux L, Pai AA, Gicquel B, Marioni JC and Gilad Y. Deciphering the genetic architecture of variation in the immune response to Mycobacterium tuberculosis infection. *Proc Natl Acad Sci U S A*. 2012;109:1204-9.
51. Lee MN, Ye C, Villani AC, Raj T, Li W, Eisenhaure TM, Imboywa SH, Chipendo PI, Ran FA, Slowikowski K, Ward LD, Raddassi K, McCabe C, Lee MH, Frohlich IY, Hafler DA, Kellis M, Raychaudhuri S, Zhang F, Stranger BE, Benoist CO, De Jager PL, Regev A and Hacohen N. Common genetic variants modulate pathogen-sensing responses in human dendritic cells. *Science*. 2014;343:1246980.
52. Fischer D, Wahlfors T, Mattila H, Oja H, Tammela TL and Schleutker J. MiRNA Profiles in Lymphoblastoid Cell Lines of Finnish Prostate Cancer Families. *PLoS One*. 2015;10:e0127427.
53. Huang RS, Gamazon ER, Ziliak D, Wen Y, Im HK, Zhang W, Wing C, Duan S, Bleibel WK, Cox NJ and Dolan ME. Population differences in microRNA expression and biological implications. *RNA Biol*. 2011;8:692-701.
54. Degner JF, Pai AA, Pique-Regi R, Veyrieras JB, Gaffney DJ, Pickrell JK, De Leon S, Michelini K, Lewellen N, Crawford GE, Stephens M, Gilad Y and Pritchard JK. DNase I sensitivity QTLs are a major determinant of human expression variation. *Nature*. 2012;482:390-4.
55. del Rosario RC, Poschmann J, Rouam SL, Png E, Khor CC, Hibberd ML and Prabhakar S. Sensitive detection of chromatin-altering polymorphisms reveals autoimmune disease mechanisms. *Nat Methods*. 2015;12:458-64.
56. Battle A, Khan Z, Wang SH, Mitrano A, Ford MJ, Pritchard JK and Gilad Y. Genomic variation. Impact of regulatory variation from RNA to protein. *Science*. 2015;347:664-7.
57. Zhang X, Joehanes R, Chen BH, Huan T, Ying S, Munson PJ, Johnson AD, Levy D and O'Donnell CJ. Identification of common genetic variants controlling transcript isoform variation in human whole blood. *Nat Genet*. 2015;47:345-52.
58. Huan T, Rong J, Liu C, Zhang X, Tanriverdi K, Joehanes R, Chen BH, Murabito JM, Yao C, Courchesne P, Munson PJ, O'Donnell CJ, Cox N, Johnson AD, Larson MG, Levy D and Freedman JE. Genome-wide identification of microRNA expression quantitative trait loci. *Nature communications*. 2015;6:6601.
59. Greenawalt DM, Dobrin R, Chudin E, Hatoum IJ, Suver C, Beaulaurier J, Zhang B, Castro V, Zhu J, Sieberts SK, Wang S, Molony C, Heymsfield SB, Kemp DM, Reitman ML, Lum PY, Schadt EE and Kaplan LM. A survey of the genetics of stomach, liver, and adipose gene expression from a morbidly obese cohort. *Genome Res*. 2011;21:1008-16.
60. Kompass KS and Witte JS. Co-regulatory expression quantitative trait loci mapping: method and application to endometrial cancer. *BMC Med Genomics*. 2011;4:6.
61. Li Q, Seo JH, Stranger B, McKenna A, Pe'er I, Laframboise T, Brown M, Tyekucheva S and Freedman ML. Integrative eQTL-based analyses reveal the biology of breast cancer risk loci. *Cell*. 2013;152:633-41.
62. Wang X, Tang H, Teng M, Li Z, Li J, Fan J, Zhong L, Sun X, Xu J, Chen G, Chen D, Wang Z, Xing T, Zhang J, Huang L, Wang S, Peng X, Qin S, Shi Y and Peng Z. Mapping of hepatic expression quantitative trait loci (eQTLs) in a Han Chinese population. *J Med Genet*. 2014;51:319-26.
63. Schroder A, Klein K, Winter S, Schwab M, Bonin M, Zell A and Zanger UM. Genomics of ADME gene expression: mapping expression quantitative trait loci relevant

for absorption, distribution, metabolism and excretion of drugs in human liver.

*Pharmacogenomics J.* 2013;13:12-20.

64. Innocenti F, Cooper GM, Stanaway IB, Gamazon ER, Smith JD, Mirkov S, Ramirez J, Liu W, Lin YS, Moloney C, Aldred SF, Trinklein ND, Schuetz E, Nickerson DA, Thummel KE, Rieder MJ, Rettie AE, Ratain MJ, Cox NJ and Brown CD. Identification, replication, and functional fine-mapping of expression quantitative trait loci in primary human liver tissue. *PLoS Genet.* 2011;7:e1002078.

65. Schadt EE, Molony C, Chudin E, Hao K, Yang X, Lum PY, Kasarskis A, Zhang B, Wang S, Suver C, Zhu J, Millstein J, Sieberts S, Lamb J, GuhaThakurta D, Derry J, Storey JD, Avila-Campillo I, Kruger MJ, Johnson JM, Rohl CA, van Nas A, Mehrabian M, Drake TA, Lusis AJ, Smith RC, Guengerich FP, Strom SC, Schuetz E, Rushmore TH and Ulrich R. Mapping the genetic architecture of gene expression in human liver. *PLoS Biol.* 2008;6:e107.

66. Grundberg E, Kwan T, Ge B, Lam KC, Koka V, Kindmark A, Mallmin H, Dias J, Verlaan DJ, Ouimet M, Sinnett D, Rivadeneira F, Estrada K, Hofman A, van Meurs JM, Uitterlinden A, Beaulieu P, Graziani A, Harmsen E, Ljunggren O, Ohlsson C, Mellstrom D, Karlsson MK, Nilsson O and Pastinen T. Population genomics in a disease targeted primary cell model. *Genome Res.* 2009;19:1942-52.

67. Kabakchiev B and Silverberg MS. Expression quantitative trait loci analysis identifies associations between genotype and gene expression in human intestine. *Gastroenterology.* 2013;144:1488-96, 1496 e1-3.

68. Hulur I, Gamazon ER, Skol AD, Xicola RM, Llor X, Onel K, Ellis NA and Kupfer SS. Enrichment of inflammatory bowel disease and colorectal cancer risk variants in colon expression quantitative trait loci. *BMC Genomics.* 2015;16:138.

69. Ongen H, Andersen CL, Bramsen JB, Oster B, Rasmussen MH, Ferreira PG, Sandoval J, Vidal E, Whiffin N, Planchon A, Padioleau I, Bielser D, Romano L, Tomlinson I, Houlston RS, Esteller M, Orntoft TF and Dermitzakis ET. Putative cis-regulatory drivers in colorectal cancer. *Nature.* 2014;512:87-90.

70. Keildson S, Fadista J, Ladenvall C, Hedman AK, Elgzyri T, Small KS, Grundberg E, Nica AC, Glass D, Richards JB, Barrett A, Nisbet J, Zheng HF, Ronn T, Strom K, Eriksson KF, Prokopenko I, Consortium M, Consortium D, Mu TC, Spector TD, Dermitzakis ET, Deloukas P, McCarthy MI, Rung J, Groop L, Franks PW, Lindgren CM and Hansson O. Expression of phosphofructokinase in skeletal muscle is influenced by genetic variation and associated with insulin sensitivity. *Diabetes.* 2014;63:1154-65.

71. Quigley DA, Fiorito E, Nord S, Van Loo P, Alnaes GG, Fleischer T, Tost J, Moen Volla HK, Tramm T, Overgaard J, Bukholm IR, Hurtado A, Balmain A, Borresen-Dale AL and Kristensen V. The 5p12 breast cancer susceptibility locus affects MRPS30 expression in estrogen-receptor positive tumors. *Mol Oncol.* 2014;8:273-84.

72. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, Speed D, Lynch AG, Samarajiwa S, Yuan Y, Graf S, Ha G, Haffari G, Bashashati A, Russell R, McKinney S, Group M, Langerod A, Green A, Provenzano E, Wishart G, Pinder S, Watson P, Markowitz F, Murphy L, Ellis I, Purushotham A, Borresen-Dale AL, Brenton JD, Tavare S, Caldas C and Aparicio S. The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature.* 2012;486:346-52.

73. Luo W, Obeidat M, Di Narzo AF, Chen R, Sin DD, Pare PD and Hao K. Airway Epithelial Expression Quantitative Trait Loci Reveal Genes Underlying Asthma and Other Airway Diseases. *Am J Respir Cell Mol Biol.* 2016;54:177-87.



74. Gao C, Tignor NL, Salit J, Strulovici-Barel Y, Hackett NR, Crystal RG and Mezey JG. HEFT: eQTL analysis of many thousands of expressed genes while simultaneously controlling for hidden factors. *Bioinformatics*. 2014;30:369-76.
75. Lamontagne M, Couture C, Postma DS, Timens W, Sin DD, Pare PD, Hogg JC, Nickle D, Laviolette M and Bosse Y. Refining susceptibility loci of chronic obstructive pulmonary disease with lung eqtls. *PLoS One*. 2013;8:e70220.
76. Hao K, Bosse Y, Nickle DC, Pare PD, Postma DS, Laviolette M, Sandford A, Hackett TL, Daley D, Hogg JC, Elliott WM, Couture C, Lamontagne M, Brandsma CA, van den Berge M, Koppelman G, Reicin AS, Nicholson DW, Malkov V, Derry JM, Suver C, Tsou JA, Kulkarni A, Zhang C, Vessey R, Opitck GJ, Curtis SP, Timens W and Sin DD. Lung eQTLs to help reveal the molecular underpinnings of asthma. *PLoS Genet*. 2012;8:e1003029.
77. Ding J, Gudjonsson JE, Liang L, Stuart PE, Li Y, Chen W, Weichenthal M, Ellinghaus E, Franke A, Cookson W, Nair RP, Elder JT and Abecasis GR. Gene expression in skin and lymphoblastoid cells: Refined statistical method reveals extensive overlap in cis-eQTL signals. *Am J Hum Genet*. 2010;87:779-89.
78. Wagner JR, Busche S, Ge B, Kwan T, Pastinen T and Blanchette M. The relationship between DNA methylation, genetic and expression inter-individual variation in untransformed human fibroblasts. *Genome Biol*. 2014;15:R37.
79. Qiu W, Cho MH, Riley JH, Anderson WH, Singh D, Bakke P, Gulsvik A, Litonjua AA, Lomas DA, Crapo JD, Beaty TH, Celli BR, Rennard S, Tal-Singer R, Fox SM, Silverman EK, Hersh CP and Investigators E. Genetics of sputum gene expression in chronic obstructive pulmonary disease. *PLoS One*. 2011;6:e24395.
80. Fadista J, Vikman P, Laakso EO, Mollet IG, Esguerra JL, Taneera J, Storm P, Osmark P, Ladvall C, Prasad RB, Hansson KB, Finotello F, Uvebrant K, Ofori JK, Di Camillo B, Krus U, Cilio CM, Hansson O, Eliasson L, Rosengren AH, Renstrom E, Wollheim CB and Groop L. Global genomic and transcriptomic analysis of human pancreatic islets reveals novel genes influencing glucose metabolism. *Proc Natl Acad Sci U S A*. 2014;111:13924-9.
81. Larson NB, McDonnell S, French AJ, Fogarty Z, Cheville J, Middha S, Riska S, Baheti S, Nair AA, Wang L, Schaid DJ and Thibodeau SN. Comprehensively evaluating cis-regulatory variation in the human prostate transcriptome by using gene-level allele-specific expression. *Am J Hum Genet*. 2015;96:869-82.
82. Singh T, Levine AP, Smith PJ, Smith AM, Segal AW and Barrett JC. Characterization of expression quantitative trait loci in the human colon. *Inflamm Bowel Dis*. 2015;21:251-6.
83. Koopmann TT, Adriaens ME, Moerland PD, Marsman RF, Westerveld ML, Lal S, Zhang T, Simmons CQ, Bacsko I, dos Remedios C, Bishopric NH, Varro A, George AL, Jr., Lodder EM and Bezzina CR. Genome-wide identification of expression quantitative trait loci (eQTLs) in human heart. *PLoS One*. 2014;9:e97380.
84. Lin H, Dolmatova EV, Morley MP, Lunetta KL, McManus DD, Magnani JW, Margulies KB, Hakonarson H, del Monte F, Benjamin EJ, Cappola TP and Ellinor PT. Gene expression and genetic variation in human atria. *Heart Rhythm*. 2014;11:266-71.
85. Rantalainen M, Herrera BM, Nicholson G, Bowden R, Wills QF, Min JL, Neville MJ, Barrett A, Allen M, Rayner NW, Fleckner J, McCarthy MI, Zondervan KT, Karpe F, Holmes CC and Lindgren CM. MicroRNA expression in abdominal and gluteal adipose tissue is associated with mRNA expression levels and partly genetically driven. *PLoS One*. 2011;6:e27338.

86. Gamazon ER, Innocenti F, Wei R, Wang L, Zhang M, Mirkov S, Ramirez J, Huang RS, Cox NJ, Ratain MJ and Liu W. A genome-wide integrative study of microRNAs in human liver. *BMC Genomics*. 2013;14:395.
87. Olsson AH, Volkov P, Bacos K, Dayeh T, Hall E, Nilsson EA, Ladenvall C, Ronn T and Ling C. Genome-wide associations between genetic and epigenetic variation influence mRNA expression and insulin secretion in human pancreatic islets. *PLoS Genet*. 2014;10:e1004735.
88. Li Q, Stram A, Chen C, Kar S, Gayther S, Pharoah P, Haiman C, Stranger B, Kraft P and Freedman ML. Expression QTL-based analyses reveal candidate causal genes and loci across five tumor types. *Hum Mol Genet*. 2014;23:5294-302.
89. Kim Y, Xia K, Tao R, Giusti-Rodriguez P, Vladimirov V, van den Oord E and Sullivan PF. A meta-analysis of gene expression quantitative trait loci in brain. *Transl Psychiatry*. 2014;4:e459.
90. Zou F, Chai HS, Younkin CS, Allen M, Crook J, Pankratz VS, Carrasquillo MM, Rowley CN, Nair AA, Middha S, Maharjan S, Nguyen T, Ma L, Malphrus KG, Palusak R, Lincoln S, Biscoglio G, Georgescu C, Kouri N, Kolbert CP, Jen J, Haines JL, Mayeux R, Pericak-Vance MA, Farrer LA, Schellenberg GD, Alzheimer's Disease Genetics C, Petersen RC, Graff-Radford NR, Dickson DW, Younkin SG and Ertekin-Taner N. Brain expression genome-wide association study (eGWAS) identifies human disease-associated variants. *PLoS Genet*. 2012;8:e1002707.
91. Webster JA, Gibbs JR, Clarke J, Ray M, Zhang W, Holmans P, Rohrer K, Zhao A, Marlowe L, Kaleem M, McCorquodale DS, 3rd, Cuello C, Leung D, Bryden L, Nath P, Zismann VL, Joshupura K, Huentelman MJ, Hu-Lince D, Coon KD, Craig DW, Pearson JV, Group NA-N, Heward CB, Reiman EM, Stephan D, Hardy J and Myers AJ. Genetic control of human brain transcript expression in Alzheimer disease. *Am J Hum Genet*. 2009;84:445-58.
92. Ramasamy A, Trabzuni D, Guelfi S, Varghese V, Smith C, Walker R, De T, Consortium UKBE, North American Brain Expression C, Coin L, de Silva R, Cookson MR, Singleton AB, Hardy J, Ryten M and Weale ME. Genetic variability in the regulation of gene expression in ten regions of the human brain. *Nat Neurosci*. 2014;17:1418-1428.
93. Zhang B, Gaiteri C, Bodea LG, Wang Z, McElwee J, Podtelezhnikov AA, Zhang C, Xie T, Tran L, Dobrin R, Fluder E, Clurman B, Melquist S, Narayanan M, Suver C, Shah H, Mahajan M, Gillis T, Mysore J, MacDonald ME, Lamb JR, Bennett DA, Molony C, Stone DJ, Gudnason V, Myers AJ, Schadt EE, Neumann H, Zhu J and Emilsson V. Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. *Cell*. 2013;153:707-20.
94. Kim S, Cho H, Lee D and Webster MJ. Association between SNPs and gene expression in multiple regions of the human brain. *Transl Psychiatry*. 2012;2:e113.
95. Gamazon ER, Badner JA, Cheng L, Zhang C, Zhang D, Cox NJ, Gershon ES, Kelsoe JR, Greenwood TA, Nievergelt CM, Chen C, McKinney R, Shilling PD, Schork NJ, Smith EN, Bloss CS, Nurnberger JI, Edenberg HJ, Foroud T, Koller DL, Scheftner WA, Coryell W, Rice J, Lawson WB, Nwulia EA, Hipolito M, Byerley W, McMahon FJ, Schulze TG, Berrettini WH, Potash JB, Zandi PP, Mahon PB, McInnis MG, Zollner S, Zhang P, Craig DW, Szelinger S, Barrett TB and Liu C. Enrichment of cis-regulatory gene expression SNPs and methylation quantitative trait loci among bipolar disorder susceptibility variants. *Mol Psychiatry*. 2013;18:340-6.
96. Gibbs JR, van der Brug MP, Hernandez DG, Traynor BJ, Nalls MA, Lai SL, Arepalli S, Dillman A, Rafferty IP, Troncoso J, Johnson R, Zielke HR, Ferrucci L, Longo DL,

- Cookson MR and Singleton AB. Abundant quantitative trait loci exist for DNA methylation and gene expression in human brain. *PLoS Genet.* 2010;6:e1000952.
97. Shpak M, Hall AW, Goldberg MM, Derryberry DZ, Ni Y, Iyer VR and Cowperthwaite MC. An eQTL analysis of the human glioblastoma multiforme genome. *Genomics.* 2014;103:252-63.
  98. Colantuoni C, Lipska BK, Ye T, Hyde TM, Tao R, Leek JT, Colantuoni EA, Elkahoul AG, Herman MM, Weinberger DR and Kleinman JE. Temporal dynamics and genetic control of transcription in the human prefrontal cortex. *Nature.* 2011;478:519-23.
  99. Liu C, Cheng L, Badner JA, Zhang D, Craig DW, Redman M and Gershon ES. Whole-genome association mapping of gene expression in the human prefrontal cortex. *Mol Psychiatry.* 2010;15:779-84.
  100. Pers TH, Karjalainen JM, Chan Y, Westra HJ, Wood AR, Yang J, Lui JC, Vedantam S, Gustafsson S, Esko T, Frayling T, Speliotes EK, Genetic Investigation of ATC, Boehnke M, Raychaudhuri S, Fehrmann RS, Hirschhorn JN and Franke L. Biological interpretation of genome-wide association studies using predicted gene functions. *Nature communications.* 2015;6:5890.
  101. Bowden J, Davey Smith G and Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol.* 2015;44:512-25.
  102. Bowden J, Davey Smith G, Haycock PC and Burgess S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol.* 2016;40:304-14.
  103. Burgess S, Bowden J, Fall T, Ingelsson E and Thompson SG. Sensitivity Analyses for Robust Causal Inference from Mendelian Randomization Analyses with Multiple Genetic Variants. *Epidemiology.* 2017;28:30-42.
  104. Harris TB, Launer LJ, Eiriksdottir G, Kjartansson O, Jonsson PV, Sigurdsson G, Thorgeirsson G, Aspelund T, Garcia ME, Cotch MF, Hoffman HJ and Gudnason V. Age, Gene/Environment Susceptibility-Reykjavik Study: multidisciplinary applied phenomics. *Am J Epidemiol.* 2007;165:1076-87.
  105. Danesh J, Wheeler JG, Hirschfield GM, Eda S, Eiriksdottir G, Rumley A, Lowe GD, Pepys MB and Gudnason V. C-reactive protein and other circulating markers of inflammation in the prediction of coronary heart disease. *N Engl J Med.* 2004;350:1387-97.
  106. The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. The ARIC investigators. *American journal of epidemiology.* 1989;129:687-702.
  107. Folsom AR, Wu KK, Rosamond WD, Sharrett AR and Chambless LE. Prospective study of hemostatic factors and incidence of coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) Study. *Circulation.* 1997;96:1102-8.
  108. Chambless LE, McMahon R, Wu K, Folsom A, Finch A and Shen YL. Short-term intraindividual variability in hemostasis factors. The ARIC Study. Atherosclerosis Risk in Communities Intraindividual Variability Study. *Ann Epidemiol.* 1992;2:723-33.
  109. Strachan DP, Rudnicka AR, Power C, Shepherd P, Fuller E, Davis A, Gibb I, Kumari M, Rumley A, Macfarlane GJ, Rahi J, Rodgers B and Stansfeld S. Lifecourse influences on health among British adults: effects of region of residence in childhood and adulthood. *Int J Epidemiol.* 2007;36:522-31.
  110. Barrett JC, Clayton DG, Concannon P, Akolkar B, Cooper JD, Erlich HA, Julier C, Morahan G, Nerup J, Nierras C, Plagnol V, Pociot F, Schuilenburg H, Smyth DJ, Stevens H, Todd JA, Walker NM, Rich SS and Type 1 Diabetes Genetics C. Genome-wide association

- study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nat Genet.* 2009;41:703-7.
111. Moffatt MF, Gut IG, Demenais F, Strachan DP, Bouzigon E, Heath S, von Mutius E, Farrall M, Lathrop M, Cookson W and Consortium G. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med.* 2010;363:1211-1221.
  112. The Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature.* 2007;447:661-78.
  113. Rudnicka AR, Rumley A, Lowe GD and Strachan DP. Diurnal, seasonal, and blood-processing patterns in levels of circulating fibrinogen, fibrin D-dimer, C-reactive protein, tissue plasminogen activator, and von Willebrand factor in a 45-year-old population. *Circulation.* 2007;115:996-1003.
  114. Friedman GD, Cutter GR, Donahue RP, Hughes GH, Hulley SB, Jacobs DR, Jr., Liu K and Savage PJ. CARDIA: study design, recruitment, and some characteristics of the examined subjects. *Journal of clinical epidemiology.* 1988;41:1105-16.
  115. Green D, Ruth KJ, Folsom AR and Liu K. Hemostatic factors in the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Arterioscler Thromb.* 1994;14:686-93.
  116. Batlle J L-FM. Laboratory assays for von Willebrand factor. In: R. Z. Zimmerman TS, ed. *Coagulation and Bleeding Disorders* New York; 1989: 330-331.
  117. Fried LP, Borhani NO, Enright P, Furberg CD, Gardin JM, Kronmal RA, Kuller LH, Manolio TA, Mittelmark MB, Newman A and et al. The Cardiovascular Health Study: design and rationale. *Ann Epidemiol.* 1991;1:263-76.
  118. Cushman M, Cornell ES, Howard PR, Bovill EG and Tracy RP. Laboratory methods and quality assurance in the Cardiovascular Health Study. *Clin Chem.* 1995;41:264-70.
  119. Feinleib M, Kannel WB, Garrison RJ, McNamara PM and Castelli WP. The Framingham Offspring Study. Design and preliminary data. *Prev Med.* 1975;4:518-25.
  120. Tofler GH, Massaro J, Levy D, Mittleman M, Sutherland P, Lipinska I, Muller JE and D'Agostino RB. Relation of the prothrombotic state to increasing age (from the Framingham Offspring Study). *Am J Cardiol.* 2005;96:1280-3.
  121. Souto JC, Almasy L, Borrell M, Gari M, Martinez E, Mateo J, Stone WH, Blangero J and Fontcuberta J. Genetic determinants of hemostasis phenotypes in Spanish families. *Circulation.* 2000;101:1546-51.
  122. Winkelmann BR, Marz W, Boehm BO, Zotz R, Hager J, Hellstern P, Senges J and Group LS. Rationale and design of the LURIC study--a resource for functional genomics, pharmacogenomics and long-term prognosis of cardiovascular disease. *Pharmacogenomics.* 2001;2:S1-73.
  123. Oudot-Mellakh T, Cohen W, Germain M, Saut N, Kallel C, Zelenika D, Lathrop M, Tregouet DA and Morange PE. Genome wide association study for plasma levels of natural anticoagulant inhibitors and protein C anticoagulant pathway: the MARTHA project. *Br J Haematol.* 2012;157:230-9.
  124. Cohen W, Castelli C, Alessi MC, Aillaud MF, Bouvet S, Saut N, Brunet D, Barthet MC, Tregouet DA, Lavigne G and Morange PE. ABO blood group and von Willebrand factor levels partially explained the incomplete penetrance of congenital thrombophilia. *Arterioscler Thromb Vasc Biol.* 2012;32:2021-8.
  125. Blom JW, Doggen CJ, Osanto S and Rosendaal FR. Malignancies, prothrombotic mutations, and the risk of venous thrombosis. *JAMA.* 2005;293:715-22.
  126. Bild DE, Bluemke DA, Burke GL, Detrano R, Diez Roux AV, Folsom AR, Greenland P, Jacob DR, Jr., Kronmal R, Liu K, Nelson JC, O'Leary D, Saad MF, Shea S, Szklo M and

Tracy RP. Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol*. 2002;156:871-81.

127. Hofman A, Brusselle GG, Darwish Murad S, van Duijn CM, Franco OH, Goedegebure A, Ikram MA, Klaver CC, Nijsten TE, Peeters RP, Stricker BH, Tiemeier HW, Uitterlinden AG and Vernooij MW. The Rotterdam Study: 2016 objectives and design update. *European journal of epidemiology*. 2015;30:661-708.

## **STUDY SOURCES OF FUNDING**

The **Age, Gene/Environment Susceptibility (AGES)**--Reykjavik Study has been funded by NIH contract N01-AG-12100, the NIA Intramural Research Program (Z01AG007380), Hjartavernd (the Icelandic Heart Association), and the Althingi (the Icelandic Parliament). The study is approved by the Icelandic National Bioethics Committee, VSN: 00-063.

**Atherosclerosis Risk in Communities (ARIC)** study is carried out as a collaborative study supported by the National Heart, Lung, and Blood Institute (NHLBI) contracts (HHSN268201100005C, HHSN268201100006C, HHSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C), R01HL087641, R01HL59367 and R01HL086694; National Human Genome Research Institute contract U01HG004402; and National Institutes of Health contract HHSN268200625226C. The authors thank the staff and participants of the ARIC study for their important contributions.

Infrastructure was partly supported by Grant Number UL1RR025005, a component of the National Institutes of Health and NIH Roadmap for Medical Research. The authors thank the staff and participants of the ARIC study for their important contributions.

Funding support for “Building on GWAS for NHLBI-diseases: the U.S. CHARGE consortium” was provided by the NIH through the American Recovery and Reinvestment Act of 2009 (ARRA) (5RC2HL102419).

We acknowledge use of phenotype and genotype data from the **British 1958 Birth Cohort (B58C)** DNA collection, funded by the Medical Research Council grant G0000934 and the Wellcome Trust grant 068545/Z/02. Genotyping for the B58C-WTCCC subset was funded by the Wellcome Trust grant 076113/B/04/Z. The B58C-T1DGC genotyping utilized resources provided by the Type 1 Diabetes Genetics Consortium, a collaborative clinical study sponsored by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National Institute of Allergy and Infectious Diseases (NIAID), National Human Genome Research Institute (NHGRI), National Institute of Child Health and Human Development (NICHD), and Juvenile Diabetes Research Foundation International (JDRF) and supported by U01 DK062418. B58C-T1DGC GWAS data were deposited by the Diabetes and Inflammation Laboratory, Cambridge Institute for Medical Research (CIMR), University of Cambridge, which is funded by Juvenile Diabetes Research Foundation International, the Wellcome Trust and the National Institute for Health Research Cambridge Biomedical Research Centre; the CIMR is in receipt of a Wellcome Trust Strategic Award (079895). The B58C-GABRIEL genotyping was supported by a contract from the European Commission Framework Programme 6 (018996) and grants from the French Ministry of Research.

The **Coronary Artery Risk Development in Young Adults Study (CARDIA)** is conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with the University of Alabama at Birmingham (HHSN268201300025C & HSN268201300026C), Northwestern University (HHSN268201300027C), University of Minnesota (HHSN268201300028C), Kaiser Foundation Research Institute (HHSN268201300029C), and Johns Hopkins University School of Medicine

(HHSN268200900041C). CARDIA is also partially supported by the Intramural Research Program of the National Institute on Aging (NIA) and an intra-agency agreement between NIA and NHLBI (AG0005). Genotyping was funded as part of the National Heart Lung and Blood Institute Candidate-gene Association Resource (N01-HC-65226) and the NHGRI Gene Environment Association Studies (GENEVA) (U01-HG004729, U01-HG04424, and U01-HG004446). This manuscript has been reviewed and approved by CARDIA for scientific content.

**Cardiovascular Health Study (CHS)** research was supported by National Heart, Lung and Blood Institute contracts HHSN268201200036C, HHSN268200800007C, HHSN268200960009C, HHSN268201800001C, N01HC55222, N01HC85079, N01HC85080, N01HC85081, N01HC85082, N01HC85083, N01HC85086; and NHLBI grants U01HL080295, R01HL085251, R01HL087652, R01HL105756, R01HL103612, R01HL120393, and R01HL130114 with additional contribution from the National Institute of Neurological Disorders and Stroke (NINDS). Additional support was provided through R01AG023629 from the National Institute on Aging (NIA). A full list of principal CHS investigators and institutions can be found at CHS-NHLBI.org. The provision of genotyping data was supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR000124, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center.

The **CROATIA-Vis** study was funded by grants from the Medical Research Council (UK) and Republic of Croatia Ministry of Science, Education and Sports research grants to I.R. (108-1080315-0302). We would like to acknowledge the staff of several institutions in Croatia that supported the field work, including but not limited to The University of Split and Zagreb Medical Schools, the Institute for Anthropological Research in Zagreb and Croatian Institute for Public Health. The SNP genotyping for the CROATIA-Vis cohort was performed in the core genotyping laboratory of Clinical Research Facility at the Western General Hospital, Edinburgh, Scotland

**Framingham Heart Study (FHS)** was partially supported by the National Heart, Lung, and Blood Institute's (NHLBI's) Framingham Heart Study (Contract No. N01-HC-25195) and its contract with Affymetrix, Inc. for genotyping services (Contract No. N02-HL-6-4278). A portion of this research utilized the Linux Cluster for Genetic Analysis (LinGA-II) funded by the Robert Dawson Evans Endowment of the Department of Medicine at Boston University School of Medicine and Boston Medical Center. Geoffrey Tofler acknowledges funding from the National Institutes of Health (R01-HL-48157). The analyses reflect intellectual input and resource development from the Framingham Heart Study investigators participating in the SNP Health Association Resource (SHARe) project. The views expressed in this manuscript are those of the authors and do not necessarily represent the views of the National Heart, Lung, and Blood Institute; the National Institutes of Health; or the U.S. Department of Health and Human Services.

The **Genes and Blood Clotting Study (GABC)** was funded by the National Institutes of Health grants, R37HL039693 and R01HL112642. David Ginsburg is a Howard Hughes Investigator.



**Genetic Analysis for Idiopathic Thrombophilia (GAIT)** project was supported partially by grants PI-11/0184, PI-14/0582 and Red Investigación Cardiovascular RD12/0042/0032 from the Instituto Carlos III (Fondo de Investigación Sanitaria–FIS), and 2014SGR-402

Consolidated Research Group of the Generalitat de Catalunya.

**Genetic Study of Atherosclerosis Risk (GeneSTAR)** was supported by grants from the National Institutes of Health/National Heart, Lung, and Blood Institute (U01 HL72518, HL087698) and by a grant from the National Institutes of Health/National Center for Research Resources (M01-RR000052) to the Johns Hopkins General Clinical Research Center.

We thank the **LURIC study** team who were either temporarily or permanently involved in patient recruitment as well as sample and data handling, in addition to the laboratory staff at the Ludwigshafen General Hospital and the Universities of Freiburg and Ulm, Germany. LURIC was supported by the 7th Framework Program RiskyCAD (grant agreement number 305739) of the European Union. The work of W.M. and M.E.K. is supported as part of the Competence Cluster of Nutrition and Cardiovascular Health (nutriCARD) which is funded by the German Federal Ministry of Education and Research.

The **MARTHA** project was supported by grants from the Program Hospitalier de Recherche Clinique. MARTHA genetics research programs are supported and funded by the GenMed LABEX (ANR-10-LBX-0013), the ICAN Institute for Cardiometabolism

and Nutrition (ANR-10-IHU-05) and the French INvestigation Network on Venous Thrombo-Embolicism (INNOVTE). Statistical analysis of the MARTHA data were performed on the C2BIG computing cluster funded by the Région Ile de France, Pierre and Marie Curie University and the ICAN Institute for Cardiometabolism and Nutrition.

The **MEGA** study was supported by Netherlands Heart Foundation (NHS 98.113), the Dutch Cancer Foundation (RUL 99/1992), the Netherlands Organisation for Scientific Research (912-03-033| 2003), and partially by the Laboratory of Excellence in Medical Genomics (GenMed LABEX ANR-10-LABX-0013). We would like to thank all colleagues from the French Centre National de Génotypage for the genotyping contribution.

**MESA** and the MESA SHARe project are conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts HHSN268201500003I, N01-HC-95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-001079, UL1-TR-001420, UL1-TR-001881, and DK063491. Funding for SHARe genotyping was provided by NHLBI Contract N02-HL-64278. Genotyping was performed at Affymetrix (Santa Clara, California, USA) and the Broad Institute of Harvard and MIT (Boston, Massachusetts, USA) using the Affymetrix Genome-Wide Human SNP Array 6.0. The provision of genotyping data was supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and

Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center.

The **Orkney Complex Disease Study (ORCADES)** was supported by the Chief Scientist Office of the Scottish Government (CZB/4/276, CZB/4/710), the Royal Society, the MRC Human Genetics Unit, Arthritis Research UK and the European Union framework program 6 EUROSPAN project (contract no. LSHG-CT-2006-018947). DNA extractions were performed at the Wellcome Trust Clinical Research Facility in Edinburgh. We would like to acknowledge the invaluable contributions of Lorraine Anderson and the research nurses in Orkney, the administrative team in Edinburgh and the people of Orkney.

The **Prospective Study of Pravastatin in the Elderly at Risk (PROSPER)** trial was supported by an investigator-initiated grant from Bristol-Myers Squibb, USA. The study was conducted, analysed and reported independently of the company. The GWAS project PHASE has received funding from the European Union's Seventh Framework Programme (FP7/2007–2013) under grant agreement HEALTH-F2-2009-223004. A part of the genotyping was funded by The Netherlands Consortium for Healthy Ageing (NGI: 05060810). Professor Dr J.W.J. is an established clinical investigator of The Netherlands Heart Foundation (2001 D 032). This work was performed as part of an ongoing collaboration of the PROSPER study group in the universities of Leiden, Glasgow and Cork.


The generation and management of GWAS genotype data for the **Rotterdam Study (RS)** is supported by the Netherlands Organisation of Scientific Research NWO

Investments (no. 175.010.2005.011, 911-03-012). This study is funded by the Research Institute for Diseases in the Elderly (014-93-015; RIDE2), the Netherlands Genomics Initiative (NGI)/Netherlands Organisation for Scientific Research (NWO) project no. 050-060-810. The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam, Netherlands Organization for the Health Research and Development (ZonMw), the Research Institute for Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam. The authors are grateful to the study participants, the staff from the Rotterdam Study and the participating general practitioners and pharmacists. We thank Pascal Arp, Mila Jhamai, Marijn Verkerk, Lizbeth Herrera, Marjolein Peters and Carolina Medina-Gomez for their help in creating the Rotterdam Study GWAS database, and Karol Estrada and Carolina Medina-Gomez for the creation and analysis of imputed data.


The **Trinity Student Study (TSS)** was supported by the Intramural Research Programs of the National Institutes of Health, the Eunice Kennedy Shriver National Institute of Child Health and Human Development and the National Human Genome Research Institute. The principle investigators for the TSS study were Lawrence C. Brody (Genome Technology Branch, National Human Genome Research Institute, National Institutes of Health), James Mills (Eunice Kennedy Shriver National Institute of Child Health and Human Development) and Anne M. Molloy (School of Medicine, Trinity College, Dublin, Ireland) who provided Karl Desch and David Ginsburg plasma samples for the determination of VWF concentrations and de-identified individual level genotyping data which was used in this study.

The Medical Research Council (MRC) and the University of Bristol support the MRC Integrative Epidemiology Unit [MC\_UU\_12013/1, MC\_UU\_12013/9, MC\_UU\_00011/1].



Supplemental Figure S1a: Regional plot of *PDHB*, *PXK*, and *KCTD6* loci for FVIII (top) and VWF (bottom) phenotypes



Supplemental Figure S1b: Regional plot of *SLC39A8* locus for FVIII (top) and VWF (bottom) phenotypes in EU only




Supplemental Figure S1c: Regional plot of *FCHO2*, *TMEM171*, and *TNPO1* loci for FVIII (top) and VWF (bottom) phenotypes







Supplemental Figure S1d: Regional plot of *HLA* locus for FVIII (top) and VWF (bottom) phenotypes




Supplemental Figure S1e: Regional plot of *GIMAP7* and *GIMAP4* loci for FVIII (top) and VWF (bottom) phenotypes




Supplemental Figure S1f: Regional plot of *SOX17* and *RP1* loci for FVIII (top) and VWF (bottom) phenotypes




Supplemental Figure S1g: Regional plot of *OR13C5* and *NIPSNAP* loci for FVIII (top) and VWF (bottom) phenotypes



Supplemental Figure S1h: Regional plot of *DAB2IP* locus for FVIII (top) and VWF (bottom) phenotypes




Supplemental Figure S1i: Regional plot of *LINC00583* and *NFIB* loci for FVIII (top) and VWF (bottom) phenotypes






Supplemental Figure S1j: Regional plot of *ST3GAL4* locus for FVIII (top) and VWF (bottom) phenotypes




Supplemental Figure S1k: Regional plot of *C2CD4B* locus for FVIII (top) and VWF (bottom) phenotypes







Supplemental Figure S11: Regional plot of *RAB5C* and *KAT2A* loci for FVIII-VWF phenotype




Supplemental Figure S1m: Regional plot of *RPL3*, *TAB1*, *SYNGR1*, and *PDGB* loci for FVIII (top) and VWF (bottom) phenotypes





Supplemental Figure S1n: Regional plot of *ARSA* locus for FVIII-VWF phenotype




(a)



(b)



(c)



Supplemental Figure S3: Figure shows the hypothesized effect of the genes found in the present study in relation to the possible regulatory points in VWF synthesis and secretion from endothelial cells. The specific regulatory point suggestion is based on previous literature evidence. Of note, VWF clearance or regulation in platelets were not studied in our *in vitro* first-pass analyses.




Figure adapted from Xiang et al. Regulation of VWF expression and secretion in health and disease. Curr Opin Hematol 2016

## **INVENT Consortium**

Philippe Amouyel,<sup>1</sup> Mariza de Andrade,<sup>2</sup> Saonli Basu,<sup>3</sup> Claudine Berr,<sup>4</sup> Jennifer A Brody,<sup>5</sup> Daniel I Chasman,<sup>6</sup> Jean-Francois Dartigues,<sup>7</sup> Aaron R Folsom,<sup>8</sup> Marine Germain,<sup>9</sup> Hugoline de Haan,<sup>10</sup> John Heit,<sup>11</sup> Jeanine Houwing-Duitermaat,<sup>12</sup> Christopher Kabrhel,<sup>13</sup> Peter Kraft,<sup>14</sup> Grégoire Legal,<sup>15,16</sup> Sara Lindström,<sup>14</sup> Ramin Monajemi,<sup>12</sup> Pierre-Emmanuel Morange,<sup>17</sup> Bruce M Psaty,<sup>5,18</sup> Pieter H Reitsma,<sup>19</sup> Paul M Ridker,<sup>20</sup> Lynda M Rose,<sup>21</sup> Frits R Rosendaal,<sup>10</sup> Noémie Saut,<sup>17</sup> Eline Slagboom,<sup>22</sup> , David Smadja<sup>23</sup> Nicholas L Smith,<sup>18,24,25</sup> Pierre Suchon,<sup>17</sup> Weihong Tang,<sup>8</sup> Kent D Taylor,<sup>26</sup> David-Alexandre Trégouët,<sup>9</sup> Christophe Tzourio,<sup>27</sup> Marieke CH de Visser,<sup>19</sup> Astrid van Hylckama Vlieg,<sup>10</sup> Lu-Chen Weng,<sup>8</sup> Kerri L Wiggins.<sup>28</sup>

## **Affiliations**

- 1 Institut Pasteur de Lille, Université de Lille Nord de France, INSERM UMR\_S 744, Lille, France; Centre Hospitalier Régional Universitaire de Lille, Lille, France.
- 2 Division of Biomedical Statistics and Informatics Mayo Clinic, Rochester, MN, USA.
- 3 University of Minnesota, Division of Biostatistics, Minneapolis, MN, USA.
- 4 INSERM Research Unit U1061, University of Montpellier I, Montpellier, France.
- 5 Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology, and Health Services, University of Washington, Seattle, WA, USA.
- 6 Division of Preventive Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02215, USA.
- 7 INSERM Research Center U897, University of Bordeaux, Bordeaux, France.
- 8 University of Minnesota, Division of Epidemiology and Community Health Minneapolis, MN, USA.
- 9 Institut National pour la Santé et la Recherche Médicale (INSERM), Unité Mixte de Recherche en Santé (UMR\_S) 1166, F-75013, Paris, France; Sorbonne Universités, Université Pierre et Marie Curie (UPMC Univ Paris 06), UMR\_S 1166, Team Genomics & Pathophysiology of Cardiovascular Diseases, F-75013, Paris, France; Institute for Cardiometabolism and Nutrition (ICAN), F-75013, Paris, France.
- 10 Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden, The Netherlands; Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands.
- 11 Division of Cardiovascular Diseases, Mayo Clinic, Rochester, MN, USA.
- 12 Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, 2300 RC Leiden, Netherlands.
- 13 Department of Emergency Medicine, Massachusetts General Hospital, Channing Network Medicine, Harvard Medical School, Boston, MA 2114, USA.
- 14 Program in Genetic Epidemiology and Statistical Genetics, Department of Epidemiology, Harvard School of Public Health, Boston, MA 2115, USA.
- 15 Université de Brest, EA3878 and CIC1412, Brest, France.
- 16 Ottawa Hospital Research Institute at the University of Ottawa, Ottawa, ON, Canada.
- 17 Laboratory of Haematology, La Timone Hospital, F-13385, Marseille, France; INSERM, UMR\_S 1062, Nutrition Obesity and Risk of Thrombosis, F-13385, Marseille, France; Aix-Marseille University, UMR\_S 1062, Nutrition Obesity and Risk of Thrombosis, F-13385, Marseille, France.
- 18 Group Health Research Institute, Group Health Cooperative, Seattle WA 98101, USA.
- 19 Einthoven Laboratory for Experimental Vascular Medicine, Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden, 2300 RC, Netherlands.



- 20 Division of Preventive Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02215, USA.
- 21 Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA 02215, USA.
- 22 Department of Molecular Epidemiology, Leiden University Medical Center, 2300 RC Leiden, Netherlands.
- 23 Université Paris Descartes, Sorbonne Paris Cité, Paris, France; AP-HP, Hôpital Européen Georges Pompidou, Service d'hématologie Biologique, Paris, France; INSERM, UMR\_S 1140, Faculté de Pharmacie, Paris, France.
- 24 Department of Epidemiology, University of Washington, Seattle WA 98195, USA.
- 25 Seattle Epidemiologic Research and Information Center, VA Office of Research and Development, Seattle WA 98108, USA.
- 26 Los Angeles Biomedical Research Institute and Department of Pediatrics, Harbor-UCLA Medical Center, Torrance CA 90502, USA.
- 27 INSERM Research Center U897, University of Bordeaux, Bordeaux, France.
- 28 Department of Medicine, University of Washington, Seattle WA 98195, USA.

## MEGA STROKE CONSORTIUM

Rainer Malik <sup>1</sup>, Ganesh Chauhan <sup>2</sup>, Matthew Traylor <sup>3</sup>, Muralidharan Sargurupremraj <sup>4,5</sup>, Yukinori Okada <sup>6,7,8</sup>, Aniket Mishra <sup>4,5</sup>, Loes Rutten-Jacobs <sup>3</sup>, Anne-Katrin Giese <sup>9</sup>, Sander W van der Laan <sup>10</sup>, Solveig Gretarsdottir <sup>11</sup>, Christopher D Anderson <sup>12,13,14,14</sup>, Michael Chong <sup>15</sup>, Hieab HH Adams <sup>16,17</sup>, Tetsuro Ago <sup>18</sup>, Peter Almgren <sup>19</sup>, Philippe Amouyel <sup>20,21</sup>, Hakan Ay <sup>22,13</sup>, Traci M Bartz <sup>23</sup>, Oscar R Benavente <sup>24</sup>, Steve Bevan <sup>25</sup>, Giorgio B Boncoraglio <sup>26</sup>, Robert D Brown, Jr. <sup>27</sup>, Adam S Butterworth <sup>28,29</sup>, Catty Carrera <sup>30,31</sup>, Cara L Carty <sup>32,33</sup>, Daniel I Chasman <sup>34,35</sup>, Wei-Min Chen <sup>36</sup>, John W Cole <sup>37</sup>, Adolfo Correa <sup>38</sup>, Ioana Cotlarciuc <sup>39</sup>, Carlos Cruchaga <sup>40,41</sup>, John Danesh <sup>28,42,43,44</sup>, Paul IW de Bakker <sup>45,46</sup>, Anita L DeStefano <sup>47,48</sup>, Marcel den Hoed <sup>49</sup>, Qing Duan <sup>50</sup>, Stefan T Engelter <sup>51,52</sup>, Guido J Falcone <sup>53,54</sup>, Rebecca F Gottesman <sup>55</sup>, Raji P Grewal <sup>56</sup>, Vilmundur Gudnason <sup>57,58</sup>, Stefan Gustafsson <sup>59</sup>, Jeffrey Haessler <sup>60</sup>, Tamara B Harris <sup>61</sup>, Ahamad Hassan <sup>62</sup>, Aki S Havulinna <sup>63,64</sup>, Susan R Heckbert <sup>65</sup>, Elizabeth G Holliday <sup>66,67</sup>, George Howard <sup>68</sup>, Fang-Chi Hsu <sup>69</sup>, Hyacinth I Hyacinth <sup>70</sup>, M Arfan Ikram <sup>16</sup>, Erik Ingelsson <sup>71,72</sup>, Marguerite R Irvin <sup>73</sup>, Xueqiu Jian <sup>74</sup>, Jordi Jiménez-Conde <sup>75</sup>, Julie A Johnson <sup>76,77</sup>, J Wouter Jukema <sup>78</sup>, Masahiro Kanai <sup>6,7,79</sup>, Keith L Keene <sup>80,81</sup>, Brett M Kissela <sup>82</sup>, Dawn O Kleindorfer <sup>82</sup>, Charles Kooperberg <sup>60</sup>, Michiaki Kubo <sup>83</sup>, Leslie A Lange <sup>84</sup>, Carl D Langefeld <sup>85</sup>, Claudia Langenberg <sup>86</sup>, Lenore J Launer <sup>87</sup>, Jin-Moo Lee <sup>88</sup>, Robin Lemmens <sup>89,90</sup>, Didier Leys <sup>91</sup>, Cathryn M Lewis <sup>92,93</sup>, Wei-Yu Lin <sup>28,94</sup>, Arne G Lindgren <sup>95,96</sup>, Erik Lorentzen <sup>97</sup>, Patrik K Magnusson <sup>98</sup>, Jane Maguire <sup>99</sup>, Ani Manichaikul <sup>36</sup>, Patrick F McArdle <sup>100</sup>, James F Meschia <sup>101</sup>, Braxton D Mitchell <sup>100,102</sup>, Thomas H Mosley <sup>103,104</sup>, Michael A Nalls <sup>105,106</sup>, Toshiharu Ninomiya <sup>107</sup>, Martin J O'Donnell <sup>15,108</sup>, Bruce M Psaty <sup>109,110,111,112</sup>, Sara L Pulit <sup>113,45</sup>, Kristiina Rannikmäe <sup>114,115</sup>, Alexander P Reiner <sup>65,116</sup>, Kathryn M Rexrode <sup>117</sup>, Kenneth Rice <sup>118</sup>, Stephen S Rich <sup>36</sup>, Paul M Ridker <sup>34,35</sup>, Natalia S Rost <sup>9,13</sup>, Peter M Rothwell <sup>119</sup>, Jerome I Rotter <sup>120,121</sup>, Tatjana Rundek <sup>122</sup>, Ralph L Sacco <sup>122</sup>, Saori Sakaue <sup>7,123</sup>, Michele M Sale <sup>124</sup>, Veikko Salomaa <sup>63</sup>, Bishwa R Sapkota <sup>125</sup>, Reinhold Schmidt <sup>126</sup>, Carsten O Schmidt <sup>127</sup>, Ulf Schminke <sup>128</sup>, Pankaj Sharma <sup>39</sup>, Agnieszka Slowik <sup>129</sup>, Cathie LM Sudlow <sup>114,115</sup>, Christian Tanislav <sup>130</sup>, Turgut Tatlisumak <sup>131,132</sup>, Kent D Taylor <sup>120,121</sup>, Vincent NS Thijs <sup>133,134</sup>, Gudmar Thorleifsson <sup>11</sup>, Unnur Thorsteinsdottir <sup>11</sup>, Steffen Tiedt <sup>1</sup>, Stella Trompet <sup>135</sup>, Christophe Tzourio <sup>5,136,137</sup>, Cornelia M van Duijn <sup>138,139</sup>, Matthew Walters <sup>140</sup>, Nicholas J Wareham <sup>86</sup>, Sylvia Wassertheil-Smoller <sup>141</sup>, James G Wilson <sup>142</sup>, Kerri L Wiggins <sup>109</sup>, Qiong Yang <sup>47</sup>, Salim Yusuf <sup>15</sup>, Najaf Amin <sup>16</sup>, Hugo S Aparicio <sup>185,48</sup>, Donna K Arnett <sup>186</sup>, John Attia <sup>187</sup>, Alexa S Beiser <sup>47,48</sup>, Claudine Berr <sup>188</sup>, Julie E Buring <sup>34,35</sup>, Mariana Bustamante <sup>189</sup>, Valeria Caso <sup>190</sup>, Yu-Ching Cheng <sup>191</sup>, Seung Hoan Choi <sup>192,48</sup>, Ayesha Chowhan <sup>185,48</sup>, Natalia Cullell <sup>31</sup>, Jean-François Dartigues <sup>193,194</sup>, Hossein Delavaran <sup>95,96</sup>, Pilar Delgado <sup>195</sup>, Marcus Dörr <sup>196,197</sup>, Gunnar Engström <sup>19</sup>, Ian Ford <sup>198</sup>, Wander S Gurpreet <sup>199</sup>, Anders Hamsten <sup>200,201</sup>, Laura Heitsch <sup>202</sup>, Atsushi Hozawa <sup>203</sup>, Laura Ibanez <sup>204</sup>, Andreea Ilinca <sup>95,96</sup>, Martin Ingelsson <sup>205</sup>, Motoki Iwasaki <sup>206</sup>, Rebecca D Jackson <sup>207</sup>, Katarina Jood <sup>208</sup>, Pekka Jousilahti <sup>63</sup>, Sara Kaffashian <sup>4,5</sup>,

Lalit Kalra <sup>209</sup>, Masahiro Kamouchi <sup>210</sup>, Takanari Kitazono <sup>211</sup>, Olafur Kjartansson <sup>212</sup>, Manja Kloss <sup>213</sup>, Peter J Koudstaal <sup>214</sup>, Jerzy Krupinski <sup>215</sup>, Daniel L Labovitz <sup>216</sup>, Cathy C Laurie <sup>118</sup>, Christopher R Levi <sup>217</sup>, Linxin Li <sup>218</sup>, Lars Lind <sup>219</sup>, Cecilia M Lindgren <sup>220,221</sup>, Vasileios Lioutas <sup>222,48</sup>, Yong Mei Liu <sup>223</sup>, Oscar L Lopez <sup>224</sup>, Hirata Makoto <sup>225</sup>, Nicolas Martinez-Majander <sup>172</sup>, Koichi Matsuda <sup>225</sup>, Naoko Minegishi <sup>203</sup>, Joan Montaner <sup>226</sup>, Andrew P Morris <sup>227,228</sup>, Elena Muiño <sup>31</sup>, Martina Müller-Nurasyid <sup>229,230,231</sup>, Bo Norrving <sup>95,96</sup>, Soichi Ogishima <sup>203</sup>, Eugenio A Parati <sup>232</sup>, Leema Reddy Peddareddygar <sup>56</sup>, Nancy L Pedersen <sup>98,233</sup>, Joanna Pera <sup>129</sup>, Markus Perola <sup>63,234</sup>, Alessandro Pezzini <sup>235</sup>, Silvana Pileggi <sup>236</sup>, Raquel Rabionet <sup>237</sup>, Iolanda Riba-Llena <sup>30</sup>, Marta Ribasés <sup>238</sup>, Jose R Romero <sup>185,48</sup>, Jaume Roquer <sup>239,240</sup>, Anthony G Rudd <sup>241,242</sup>, Antti-Pekka Sarin <sup>243,244</sup>, Ralhan Sarju <sup>199</sup>, Chloe Sarnowski <sup>47,48</sup>, Makoto Sasaki <sup>245</sup>, Claudia L Satizabal <sup>185,48</sup>, Mamoru Satoh <sup>245</sup>, Naveed Sattar <sup>246</sup>, Norie Sawada <sup>206</sup>, Gerli Sibolt <sup>172</sup>, Ásgeir Sigurdsson <sup>247</sup>, Albert Smith <sup>248</sup>, Kenji Sobue <sup>245</sup>, Carolina Soriano- Tárrega <sup>240</sup>, Tara Stanne <sup>249</sup>, O Colin Stine <sup>250</sup>, David J Stott <sup>251</sup>, Konstantin Strauch <sup>229,252</sup>, Takako Takai <sup>203</sup>, Hideo Tanaka <sup>253,254</sup>, Kozo Tanno <sup>245</sup>, Alexander Teumer <sup>255</sup>, Liisa Tomppo <sup>172</sup>, Nuria P Torres-Aguila <sup>31</sup>, Emmanuel Touze <sup>256,257</sup>, Shoichiro Tsugane <sup>206</sup>, Andre G Uitterlinden <sup>258</sup>, Einar M Valdimarsson <sup>259</sup>, Sven J van der Lee <sup>16</sup>, Henry Völzke <sup>255</sup>, Kenji Wakai <sup>253</sup>, David Weir <sup>260</sup>, Stephen R Williams <sup>261</sup>, Charles DA Wolfe <sup>241,242</sup>, Quenna Wong <sup>118</sup>, Huichun Xu <sup>191</sup>, Taiki Yamaji <sup>206</sup>, Dharambir K Sanghera <sup>125,169,170</sup>, Olle Melander <sup>19</sup>, Christina Jern <sup>171</sup>, Daniel Strbian <sup>172,173</sup>, Israel Fernandez-Cadenas <sup>31,30</sup>, W T Longstreth, Jr <sup>174,65</sup>, Arndt Rolfs <sup>175</sup>, Jun Hata <sup>107</sup>, Daniel Woo <sup>82</sup>, Jonathan Rosand <sup>12,13,14</sup>, Guillaume Pare <sup>15</sup>, Jemma C Hopewell <sup>176</sup>, Danish Saleheen <sup>177</sup>, Kari Stefansson <sup>11,178</sup>, Bradford B Worrall <sup>179</sup>, Steven J Kittner <sup>37</sup>, Sudha Seshadri <sup>180,48</sup>, Myriam Fornage <sup>74,181</sup>, Hugh S Markus <sup>3</sup>, Joanna MM Howson <sup>28</sup>, Yoichiro Kamatani <sup>6,182</sup>, Stephanie Debette <sup>4,5</sup>, Martin Dichgans <sup>1,183,184</sup>

1 Institute for Stroke and Dementia Research (ISD), University Hospital, LMU Munich, Munich, Germany

2 Centre for Brain Research, Indian Institute of Science, Bangalore, India

3 Stroke Research Group, Division of Clinical Neurosciences, University of Cambridge, UK

4 INSERM U1219 Bordeaux Population Health Research Center, Bordeaux, France

5 University of Bordeaux, Bordeaux, France

6 Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan

7 Department of Statistical Genetics, Osaka University Graduate School of Medicine, Osaka, Japan

8 Laboratory of Statistical Immunology, Immunology Frontier Research Center (WPI-IFReC), Osaka University, Suita, Japan.

- 9 Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA
- 10 Laboratory of Experimental Cardiology, Division of Heart and Lungs, University Medical Center Utrecht, University of Utrecht, Utrecht, Netherlands
- 11 deCODE genetics/AMGEN inc, Reykjavik, Iceland
- 12 Center for Genomic Medicine, Massachusetts General Hospital (MGH), Boston, MA, USA
- 13 J. Philip Kistler Stroke Research Center, Department of Neurology, MGH, Boston, MA, USA
- 14 Program in Medical and Population Genetics, Broad Institute, Cambridge, MA, USA
- 15 Population Health Research Institute, McMaster University, Hamilton, Canada
- 16 Department of Epidemiology, Erasmus University Medical Center, Rotterdam, Netherlands
- 17 Department of Radiology and Nuclear Medicine, Erasmus University Medical Center, Rotterdam, Netherlands
- 18 Department of Medicine and Clinical Science, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan
- 19 Department of Clinical Sciences, Lund University, Malmö, Sweden
- 20 Univ. Lille, Inserm, Institut Pasteur de Lille, LabEx DISTALZ-UMR1167, Risk factors and molecular determinants of aging-related diseases, F-59000 Lille, France
- 21 Centre Hosp. Univ Lille, Epidemiology and Public Health Department, F-59000 Lille, France
- 22 AA Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA
- 23 Cardiovascular Health Research Unit, Departments of Biostatistics and Medicine, University of Washington, Seattle, WA, USA
- 24 Division of Neurology, Faculty of Medicine, Brain Research Center, University of British Columbia, Vancouver, Canada
- 25 School of Life Science, University of Lincoln, Lincoln, UK
- 26 Department of Cerebrovascular Diseases, Fondazione IRCCS Istituto Neurologico "Carlo Besta", Milano, Italy
- 27 Department of Neurology, Mayo Clinic Rochester, Rochester, MN, USA

28 MRC/BHF Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

29 The National Institute for Health Research Blood and Transplant Research Unit in Donor Health and Genomics, University of Cambridge, UK

30 Neurovascular Research Laboratory, Vall d'Hebron Institut of Research, Neurology and Medicine Departments-Universitat Autònoma de Barcelona, Vall d'Hebrón Hospital, Barcelona, Spain

31 Stroke Pharmacogenomics and Genetics, Fundacio Docència i Recerca Mútua Terrassa, Terrassa, Spain

32 Children's Research Institute, Children's National Medical Center, Washington, DC, USA

33 Center for Translational Science, George Washington University, Washington, DC, USA

34 Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA, USA

35 Harvard Medical School, Boston, MA, USA

36 Center for Public Health Genomics, Department of Public Health Sciences, University of Virginia, Charlottesville, VA, USA

37 Department of Neurology, University of Maryland School of Medicine and Baltimore VAMC, Baltimore, MD, USA

38 Departments of Medicine, Pediatrics and Population Health Science, University of Mississippi Medical Center, Jackson, MS, USA

39 Institute of Cardiovascular Research, Royal Holloway University of London, UK & Ashford and St Peters Hospital, Surrey UK

40 Department of Psychiatry, The Hope Center Program on Protein Aggregation and Neurodegeneration (HPAN), Washington University, School of Medicine, St. Louis, MO, USA

41 Department of Developmental Biology, Washington University School of Medicine, St. Louis, MO, USA

42 NIHR Blood and Transplant Research Unit in Donor Health and Genomics, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

43 Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, UK

- 44 British Heart Foundation, Cambridge Centre of Excellence, Department of Medicine, University of Cambridge, Cambridge, UK
- 45 Department of Medical Genetics, University Medical Center Utrecht, Utrecht, Netherlands
- 46 Department of Epidemiology, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, Netherlands
- 47 Boston University School of Public Health, Boston, MA, USA
- 48 Framingham Heart Study, Framingham, MA, USA
- 49 Department of Immunology, Genetics and Pathology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden
- 50 Department of Genetics, University of North Carolina, Chapel Hill, NC, USA
- 51 Department of Neurology and Stroke Center, Basel University Hospital, Switzerland
- 52 Neurorehabilitation Unit, University and University Center for Medicine of Aging and Rehabilitation Basel, Felix Platter Hospital, Basel, Switzerland
- 53 Department of Neurology, Yale University School of Medicine, New Haven, CT, USA
- 54 Program in Medical and Population Genetics, The Broad Institute of Harvard and MIT, Cambridge, MA, USA
- 55 Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA
- 56 Neuroscience Institute, SF Medical Center, Trenton, NJ, USA
- 57 Icelandic Heart Association Research Institute, Kopavogur, Iceland
- 58 University of Iceland, Faculty of Medicine, Reykjavik, Iceland
- 59 Department of Medical Sciences, Molecular Epidemiology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden
- 60 Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA
- 61 Laboratory of Epidemiology and Population Science, National Institute on Aging, National Institutes of Health, Bethesda, MD, USA
- 62 Department of Neurology, Leeds General Infirmary, Leeds Teaching Hospitals NHS Trust, Leeds, UK
- 63 National Institute for Health and Welfare, Helsinki, Finland
- 64 FIMM - Institute for Molecular Medicine Finland, Helsinki, Finland

- 65 Department of Epidemiology, University of Washington, Seattle, WA, USA
- 66 Public Health Stream, Hunter Medical Research Institute, New Lambton, Australia
- 67 Faculty of Health and Medicine, University of Newcastle, Newcastle, Australia
- 68 School of Public Health, University of Alabama at Birmingham, Birmingham, AL, USA
- 69 Department of Biostatistical Sciences, Wake Forest School of Medicine, Winston-Salem, NC, USA
- 70 Aflac Cancer and Blood Disorder Center, Department of Pediatrics, Emory University School of Medicine, Atlanta, GA, USA
- 71 Department of Medicine, Division of Cardiovascular Medicine, Stanford University School of Medicine, CA, USA
- 72 Department of Medical Sciences, Molecular Epidemiology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden
- 73 Epidemiology, School of Public Health, University of Alabama at Birmingham, USA
- 74 Brown Foundation Institute of Molecular Medicine, University of Texas Health Science Center at Houston, Houston, TX, USA
- 75 Neurovascular Research Group (NEUVAS), Neurology Department, Institut Hospital del Mar d'Investigació Mèdica, Universitat Autònoma de Barcelona, Barcelona, Spain
- 76 Department of Pharmacotherapy and Translational Research and Center for Pharmacogenomics, University of Florida, College of Pharmacy, Gainesville, FL, USA
- 77 Division of Cardiovascular Medicine, College of Medicine, University of Florida, Gainesville, FL, USA
- 78 Department of Cardiology, Leiden University Medical Center, Leiden, the Netherlands
- 79 Program in Bioinformatics and Integrative Genomics, Harvard Medical School, Boston, MA, USA
- 80 Department of Biology, East Carolina University, Greenville, NC, USA
- 81 Center for Health Disparities, East Carolina University, Greenville, NC, USA
- 82 University of Cincinnati College of Medicine, Cincinnati, OH, USA
- 83 RIKEN Center for Integrative Medical Sciences, Yokohama, Japan
- 84 Department of Medicine, University of Colorado Denver, Anschutz Medical Campus, Aurora, CO, USA

85 Center for Public Health Genomics and Department of Biostatistical Sciences, Wake Forest School of Medicine, Winston-Salem, NC, USA

86 MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Institute of Metabolic Science, Cambridge Biomedical Campus, Cambridge, UK

87 Intramural Research Program, National Institute on Aging, National Institutes of Health, Bethesda, MD, USA

88 Department of Neurology, Radiology, and Biomedical Engineering, Washington University School of Medicine, St. Louis, MO, USA

89 KU Leuven – University of Leuven, Department of Neurosciences, Experimental Neurology, Leuven, Belgium

90 VIB Center for Brain & Disease Research, University Hospitals Leuven, Department of Neurology, Leuven, Belgium

91 Univ.-Lille, INSERM U 1171. CHU Lille. Lille, France

92 Department of Medical and Molecular Genetics, King's College London, London, UK

93 SGDP Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

94 Northern Institute for Cancer Research, Paul O'Gorman Building, Newcastle University, Newcastle, UK

95 Department of Clinical Sciences Lund, Neurology, Lund University, Lund, Sweden

96 Department of Neurology and Rehabilitation Medicine, Skåne University Hospital, Lund, Sweden

97 Bioinformatics Core Facility, University of Gothenburg, Gothenburg, Sweden

98 Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

99 University of Technology Sydney, Faculty of Health, Ultimo, Australia

100 Department of Medicine, University of Maryland School of Medicine, MD, USA

101 Department of Neurology, Mayo Clinic, Jacksonville, FL, USA

102 Geriatrics Research and Education Clinical Center, Baltimore Veterans Administration Medical Center, Baltimore, MD, USA

103 Division of Geriatrics, School of Medicine, University of Mississippi Medical Center, Jackson, MS, USA



- 104 Memory Impairment and Neurodegenerative Dementia Center, University of Mississippi Medical Center, Jackson, MS, USA
- 105 Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, MD, USA
- 106 Data Tecnica International, Glen Echo MD, USA
- 107 Department of Epidemiology and Public Health, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan
- 108 Clinical Research Facility, Department of Medicine, NUI Galway, Galway, Ireland
- 109 Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle, WA, USA
- 110 Department of Epidemiology, University of Washington, Seattle, WA
- 111 Department of Health Services, University of Washington, Seattle, WA, USA
- 112 Kaiser Permanente Washington Health Research Institute, Seattle, WA, USA
- 113 Brain Center Rudolf Magnus, Department of Neurology, University Medical Center Utrecht, Utrecht, The Netherlands
- 114 Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, UK
- 115 Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK
- 116 Fred Hutchinson Cancer Research Center, University of Washington, Seattle, WA, USA
- 117 Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA
- 118 Department of Biostatistics, University of Washington, Seattle, WA, USA
- 119 Nuffield Department of Clinical Neurosciences, University of Oxford, UK
- 120 Institute for Translational Genomics and Population Sciences, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, CA, USA
- 121 Division of Genomic Outcomes, Department of Pediatrics, Harbor-UCLA Medical Center, Torrance, CA, USA
- 122 Department of Neurology, Miller School of Medicine, University of Miami, Miami, FL, USA
- 123 Department of Allergy and Rheumatology, Graduate School of Medicine, the University of Tokyo, Tokyo, Japan
- 124 Center for Public Health Genomics, University of Virginia, Charlottesville, VA, USA

- 125 Department of Pediatrics, College of Medicine, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA
- 126 Department of Neurology, Medical University of Graz, Graz, Austria
- 127 University Medicine Greifswald, Institute for Community Medicine, SHIP-KEF, Greifswald, Germany
- 128 University Medicine Greifswald, Department of Neurology, Greifswald, Germany
- 129 Department of Neurology, Jagiellonian University, Krakow, Poland
- 130 Department of Neurology, Justus Liebig University, Giessen, Germany
- 131 Department of Clinical Neurosciences/Neurology, Institute of Neuroscience and Physiology, Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden
- 132 Sahlgrenska University Hospital, Gothenburg, Sweden
- 133 Stroke Division, Florey Institute of Neuroscience and Mental Health, University of Melbourne, Heidelberg, Australia
- 134 Austin Health, Department of Neurology, Heidelberg, Australia
- 135 Department of Internal Medicine, Section Gerontology and Geriatrics, Leiden University Medical Center, Leiden, the Netherlands
- 136 INSERM U1219, Bordeaux, France
- 137 Department of Public Health, Bordeaux University Hospital, Bordeaux, France
- 138 Genetic Epidemiology Unit, Department of Epidemiology, Erasmus University Medical Center Rotterdam, Netherlands
- 139 Center for Medical Systems Biology, Leiden, Netherlands
- 140 School of Medicine, Dentistry and Nursing at the University of Glasgow, Glasgow, UK
- 141 Department of Epidemiology and Population Health, Albert Einstein College of Medicine, NY, USA
- 142 Department of Physiology and Biophysics, University of Mississippi Medical Center, Jackson, MS, USA
- 143 A full list of members and affiliations appears in the Supplementary Note
- 144 Department of Human Genetics, McGill University, Montreal, Canada
- 145 Department of Pathophysiology, Institute of Biomedicine and Translation Medicine, University of Tartu, Tartu, Estonia
- 146 Department of Cardiac Surgery, Tartu University Hospital, Tartu, Estonia

- 147 Clinical Gene Networks AB, Stockholm, Sweden
- 148 Department of Genetics and Genomic Sciences, The Icahn Institute for Genomics and Multiscale Biology Icahn School of Medicine at Mount Sinai, New York, NY , USA
- 149 Department of Pathophysiology, Institute of Biomedicine and Translation Medicine, University of Tartu, Biomeedikum, Tartu, Estonia
- 150 Integrated Cardio Metabolic Centre, Department of Medicine, Karolinska Institutet, Karolinska Universitetssjukhuset, Huddinge, Sweden.
- 151 Clinical Gene Networks AB, Stockholm, Sweden
- 152 Sorbonne Universités, UPMC Univ. Paris 06, INSERM, UMR\_S 1166, Team Genomics & Pathophysiology of Cardiovascular Diseases, Paris, France
- 153 ICAN Institute for Cardiometabolism and Nutrition, Paris, France
- 154 Department of Biomedical Engineering, University of Virginia, Charlottesville, VA, USA
- 155 Group Health Research Institute, Group Health Cooperative, Seattle, WA, USA
- 156 Seattle Epidemiologic Research and Information Center, VA Office of Research and Development, Seattle, WA, USA
- 157 Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA, USA
- 158 Department of Medical Research, Bærum Hospital, Vestre Viken Hospital Trust , Gjettem, Norway
- 159 Saw Swee Hock School of Public Health, National University of Singapore and National University Health System, Singapore
- 160 National Heart and Lung Institute, Imperial College London, London, UK
- 161 Department of Gene Diagnostics and Therapeutics, Research Institute, National Center for Global Health and Medicine, Tokyo, Japan
- 162 Department of Epidemiology, Tulane University School of Public Health and Tropical Medicine, New Orleans, LA, USA
- 163 Department of Cardiology, University Medical Center Groningen, University of Groningen, Netherlands
- 164 MRC-PHE Centre for Environment and Health, School of Public Health, Department of Epidemiology and Biostatistics, Imperial College London, London, UK

- 165 Department of Epidemiology and Biostatistics, Imperial College London, London, UK
- 166 Department of Cardiology, Ealing Hospital NHS Trust, Southall, UK
- 167 National Heart, Lung and Blood Research Institute, Division of Intramural Research, Population Sciences Branch, Framingham, MA, USA
- 168 A full list of members and affiliations appears at the end of the manuscript
- 169 Department of Pharmaceutical Sciences, College of Pharmacy, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA
- 170 Oklahoma Center for Neuroscience, Oklahoma City, OK, USA
- 171 Department of Pathology and Genetics, Institute of Biomedicine, The Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden
- 172 Department of Neurology, Helsinki University Hospital, Helsinki, Finland
- 173 Clinical Neurosciences, Neurology, University of Helsinki, Helsinki, Finland
- 174 Department of Neurology, University of Washington, Seattle, WA, USA
- 175 Albrecht Kossel Institute, University Clinic of Rostock, Rostock, Germany
- 176 Clinical Trial Service Unit and Epidemiological Studies Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK
- 177 Department of Genetics, Perelman School of Medicine, University of Pennsylvania, PA, USA
- 178 Faculty of Medicine, University of Iceland, Reykjavik, Iceland
- 179 Departments of Neurology and Public Health Sciences, University of Virginia School of Medicine, Charlottesville, VA, USA
- 180 Department of Neurology, Boston University School of Medicine, Boston, MA, USA
- 181 Human Genetics Center, University of Texas Health Science Center at Houston, Houston, TX, USA
- 182 Center for Genomic Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan
- 183 Munich Cluster for Systems Neurology (SyNergy), Munich, Germany
- 184 German Center for Neurodegenerative Diseases (DZNE), Munich, Germany
- 185 Boston University School of Medicine, Boston, MA, USA
- 186 University of Kentucky College of Public Health, Lexington, KY, USA

- 187 University of Newcastle and Hunter Medical Research Institute, New Lambton, Australia
- 188 Univ. Montpellier, Inserm, U1061, Montpellier, France
- 189 Centre for Research in Environmental Epidemiology, Barcelona, Spain
- 190 Department of Neurology, Università degli Studi di Perugia, Umbria, Italy
- 191 Department of Medicine, University of Maryland School of Medicine, Baltimore, MD, USA
- 192 Broad Institute, Cambridge, MA, USA
- 193 Univ. Bordeaux, Inserm, Bordeaux Population Health Research Center, UMR 1219, Bordeaux, France
- 194 Bordeaux University Hospital, Department of Neurology, Memory Clinic, Bordeaux, France
- 195 Neurovascular Research Laboratory. Vall d'Hebron Institut of Research, Neurology and Medicine Departments-Universitat Autònoma de Barcelona. Vall d'Hebrón Hospital, Barcelona, Spain
- 196 University Medicine Greifswald, Department of Internal Medicine B, Greifswald, Germany
- 197 DZHK, Greifswald, Germany
- 198 Robertson Center for Biostatistics, University of Glasgow, Glasgow, UK
- 199 Hero DMC Heart Institute, Dayanand Medical College & Hospital, Ludhiana, India
- 200 Atherosclerosis Research Unit, Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden
- 201 Karolinska Institutet, Stockholm, Sweden
- 202 Division of Emergency Medicine, and Department of Neurology, Washington University School of Medicine, St. Louis, MO, USA
- 203 Tohoku Medical Megabank Organization, Sendai, Japan
- 204 Department of Psychiatry, Washington University School of Medicine, St. Louis, MO, USA
- 205 Department of Public Health and Caring Sciences / Geriatrics, Uppsala University, Uppsala, Sweden
- 206 Epidemiology and Prevention Group, Center for Public Health Sciences, National Cancer Center, Tokyo, Japan
- 207 Department of Internal Medicine and the Center for Clinical and Translational Science, The

Ohio State University, Columbus, OH, USA

208 Institute of Neuroscience and Physiology, the Sahlgrenska Academy at University of Gothenburg, Goteborg, Sweden

209 Department of Basic and Clinical Neurosciences, King's College London, London, UK

210 Department of Health Care Administration and Management, Graduate School of Medical Sciences, Kyushu University, Japan

211 Department of Medicine and Clinical Science, Graduate School of Medical Sciences, Kyushu University, Japan

212 Landspítali National University Hospital, Departments of Neurology & Radiology, Reykjavik, Iceland

213 Department of Neurology, Heidelberg University Hospital, Germany

214 Department of Neurology, Erasmus University Medical Center

215 Hospital Universitari Mutua Terrassa, Terrassa (Barcelona), Spain

216 Albert Einstein College of Medicine, Montefiore Medical Center, New York, NY, USA

217 John Hunter Hospital, Hunter Medical Research Institute and University of Newcastle, Newcastle, NSW, Australia

218 Centre for Prevention of Stroke and Dementia, Nuffield Department of Clinical Neurosciences, University of Oxford, UK

219 Department of Medical Sciences, Uppsala University, Uppsala, Sweden

220 Genetic and Genomic Epidemiology Unit, Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK

221 The Wellcome Trust Centre for Human Genetics, Oxford, UK

222 Beth Israel Deaconess Medical Center, Boston, MA, USA

223 Wake Forest School of Medicine, Wake Forest, NC, USA

224 Department of Neurology, University of Pittsburgh, Pittsburgh, PA, USA

225 BioBank Japan, Laboratory of Clinical Sequencing, Department of Computational biology and medical Sciences, Graduate school of Frontier Sciences, The University of Tokyo, Tokyo, Japan

226 Neurovascular Research Laboratory, Vall d'Hebron Institut of Research, Neurology and Medicine Departments-Universitat Autònoma de Barcelona. Vall d'Hebrón Hospital, Barcelona, Spain

- 227 Department of Biostatistics, University of Liverpool, Liverpool, UK
- 228 Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK
- 229 Institute of Genetic Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany
- 230 Department of Medicine I, Ludwig-Maximilians-Universität, Munich, Germany
- 231 DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, Munich, Germany
- 232 Department of Cerebrovascular Diseases, Fondazione IRCCS Istituto Neurologico "Carlo Besta", Milano, Italy
- 233 Karolinska Institutet, MEB, Stockholm, Sweden
- 234 University of Tartu, Estonian Genome Center, Tartu, Estonia, Tartu, Estonia
- 235 Department of Clinical and Experimental Sciences, Neurology Clinic, University of Brescia, Italy
- 236 Translational Genomics Unit, Department of Oncology, IRCCS Istituto di Ricerche Farmacologiche Mario Negri, Milano, Italy
- 237 Department of Genetics, Microbiology and Statistics, University of Barcelona, Barcelona, Spain
- 238 Psychiatric Genetics Unit, Group of Psychiatry, Mental Health and Addictions, Vall d'Hebron Research Institute (VHIR), Universitat Autònoma de Barcelona, Biomedical Network Research Centre on Mental Health (CIBERSAM), Barcelona, Spain
- 239 Department of Neurology, IMIM-Hospital del Mar, and Universitat Autònoma de Barcelona, Spain
- 240 IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain
- 241 National Institute for Health Research Comprehensive Biomedical Research Centre, Guy's & St. Thomas' NHS Foundation Trust and King's College London, London, UK
- 242 Division of Health and Social Care Research, King's College London, London, UK
- 243 FIMM-Institute for Molecular Medicine Finland, Helsinki, Finland
- 244 THL-National Institute for Health and Welfare, Helsinki, Finland
- 245 Iwate Tohoku Medical Megabank Organization, Iwate Medical University, Iwate, Japan
- 246 BHF Glasgow Cardiovascular Research Centre, Faculty of Medicine, Glasgow, UK
- 247 deCODE Genetics/Amgen, Inc., Reykjavik, Iceland
- 248 Icelandic Heart Association, Reykjavik, Iceland

249 Institute of Biomedicine, the Sahlgrenska Academy at University of Gothenburg, Goteborg, Sweden

250 Department of Epidemiology, University of Maryland School of Medicine, Baltimore, MD, USA

251 Institute of Cardiovascular and Medical Sciences, Faculty of Medicine, University of Glasgow, Glasgow, UK

252 Chair of Genetic Epidemiology, IBE, Faculty of Medicine, LMU Munich, Germany

253 Division of Epidemiology and Prevention, Aichi Cancer Center Research Institute, Nagoya, Japan

254 Department of Epidemiology, Nagoya University Graduate School of Medicine, Nagoya, Japan

255 University Medicine Greifswald, Institute for Community Medicine, SHIP-KEF, Greifswald, Germany

256 Department of Neurology, Caen University Hospital, Caen, France

257 University of Caen Normandy, Caen, France

258 Department of Internal Medicine, Erasmus University Medical Center, Rotterdam, Netherlands

259 Landspítali University Hospital, Reykjavik, Iceland

260 Survey Research Center, University of Michigan, Ann Arbor, MI, USA

261 University of Virginia Department of Neurology, Charlottesville, VA, USA

## **MEGASTROKE Acknowledgments**

### **METASTROKE**

*ASGC*: Australian population control data were derived from the Hunter Community Study. We also thank the University of Newcastle for funding and the men and women of the Hunter region who participated in this study. This research was funded by grants from the Australian National and Medical Health Research Council (NHMRC Project Grant ID: 569257), the Australian National Heart Foundation (NHF Project Grant ID: G 04S 1623), the University of Newcastle, the Gladys M Brawn Fellowship scheme, and the Vincent Fairfax Family Foundation in Australia. Elizabeth G Holliday was supported by a Fellowship from the National Heart Foundation and National Stroke Foundation of Australia (ID: 100071).

*BRAINS*: Bio-Repository of DNA in Stroke (BRAINS) is partly funded by a Senior Fellowship from the Department of Health (UK) to P Sharma, the Henry Smith Charity and the UK-India Education Research Institute (UKIERI) from the British Council.

*GEOS*: Genetics of Early Onset Stroke (GEOS) Study, Baltimore, USA was supported by the NIH Genes, Environment and Health Initiative (GEI) Grant U01 HG004436, as part of the GENEVA consortium under GEI, with additional support provided by the Mid-Atlantic



Nutrition and Obesity Research Center (P30 DK072488), and the Office of Research and Development, Medical Research Service, and the Baltimore Geriatrics Research, Education, and Clinical Center of the Department of Veterans Affairs. Genotyping services were provided by the Johns Hopkins University Center for Inherited Disease Research (CIDR), which is fully funded through a federal contract from the NIH to the Johns Hopkins University (contract number HHSN268200782096C). Assistance with data cleaning was provided by the GENEVA Coordinating Center (U01 HG 004446; PI Bruce S Weir). Study recruitment and assembly of datasets were supported by a Cooperative Agreement with the Division of Adult and Community Health, Centers for Disease Control and Prevention and by grants from NINDS and the NIH Office of Research on Women's Health (R01 NS45012, U01 NS069208-01).

*HPS*: Heart Protection Study (HPS) (ISRCTN48489393) was supported by the UK Medical Research Council (MRC), British Heart Foundation, Merck and Co (manufacturers of simvastatin), and Roche Vitamins Ltd (manufacturers of vitamins). Genotyping was supported by a grant to Oxford University and CNG from Merck and Co. Jemma C Hopewell acknowledges support from the British Heart Foundation (FS/14/55/30806).

*ISGS and SWISS*: The Ischemic Stroke Genetics Study (ISGS) was supported by the NINDS (R01 NS42733; PI Dr Meschia). The Sibling with Ischemic Stroke Study (SWISS) was supported by the NINDS (R01 NS39987; PI Dr Meschia). Both SWISS and ISGS received additional support, in part, from the Intramural Research Program of the National Institute on Aging (Z01 AG000954-06; PI Andrew Singleton). SWISS and ISGS used samples and clinical data from the NIH-NINDS Human Genetics Resource Center DNA and Cell Line Repository (<http://ccr.coriell.org/ninds>), human subject protocol Nos. 2003-081 and 2004-147. SWISS and ISGS used stroke-free participants from the Baltimore Longitudinal Study of Aging (BLSA) as controls with the permission of Dr Luigi Ferrucci. The inclusion of BLSA samples was supported, in part, by the Intramural Research Program of the National Institute on Aging (Z01 AG000015-50), human subject protocol No. 2003-078. This study used the high-performance computational capabilities of the Biowulf Linux cluster at the NIH (<http://biowulf.nih.gov>). For SWISS and ISGS cases of African ancestry, a subset of the Healthy Aging

in Neighborhoods of Diversity across the Life Span study (HANDLS) were used as stroke-free controls. HANDLS is funded by the National Institute of Aging (1Z01AG000513; PI Michele K. Evans).

*MGH-GASROS*: MGH Genes Affecting Stroke Risk and Outcome Study (MGH-GASROS) was supported by NINDS (U01 NS069208), the American Heart Association/Bugher Foundation Centers for Stroke Prevention Research 0775010N, the NIH and NHLBI's STAMPEED genomics research program (R01 HL087676), and a grant from the National Center for Research Resources. The Broad Institute Center for Genotyping and Analysis is supported by grant U54 RR020278 from the National Center for Research resources.

*MILANO*: MILANO: Milano - Besta Stroke Register Collection and genotyping of the Milan cases within CEDIR were supported by the Italian Ministry of Health (Grant Numbers: RC 2007/LR6, RC 2008/LR6; RC 2009/LR8; RC 2010/LR8; GR-2011-02347041). FP6 LSHM-CT-2007-037273 for the PROCARDIS control samples.

*WTCCC2-UK*: Wellcome Trust Case-Control Consortium 2 (WTCCC2) was principally funded by the Wellcome Trust, as part of the Wellcome Trust Case Control Consortium 2 project (085475/B/08/Z and

085475/Z/08/Z and WT084724MA). The Stroke Association provided additional support for collection of some of the St George's, London cases. The Oxford cases were collected as part of the Oxford Vascular Study which is funded by the MRC, Stroke Association, Dunhill Medical Trust, National Institute of Health Research (NIHR) and the NIHR Biomedical Research Centre, Oxford. The Edinburgh Stroke Study was supported by the Wellcome Trust (clinician scientist award to C Sudlow), and the Binks Trust. Sample processing occurred in the Genetics Core Laboratory of the Wellcome Trust Clinical Research Facility, Western General Hospital, Edinburgh. Much of the neuroimaging occurred in the Scottish Funding Council Brain Imaging Research Centre ([www.sbirc.ed.ac.uk](http://www.sbirc.ed.ac.uk)), Division of Clinical Neurosciences, University of Edinburgh, a core area of the Wellcome Trust Clinical Research Facility and part of the SINAPSE (Scottish Imaging Network—A Platform for Scientific Excellence) collaboration ([www.sinapse.ac.uk](http://www.sinapse.ac.uk)), funded by the Scottish Funding Council and the Chief Scientist Office. Collection of the Munich cases and data analysis was supported by the Vascular Dementia Research Foundation. M Farrall and A Helgadottir acknowledge support from the BHF Centre of Research Excellence in Oxford and the Wellcome Trust core award (090532/Z/09/Z).

*WTCCC2-D*: This project has received funding from the European Union's Horizon 2020 research and

innovation programme under grant agreements No 666881, SVDs@target (to M Dichgans) and No

667375, CoSTREAM (to M Dichgans); the DFG as part of the Munich Cluster for Systems Neurology (EXC 1010 SyNergy) and the CRC 1123 (B3)(to M Dichgans); the Corona Foundation (to M Dichgans); the Fondation Leducq (Transatlantic Network of Excellence on the Pathogenesis of Small Vessel Disease of the Brain)(to M Dichgans); the e:Med program (e:AtheroSysMed) (to M Dichgans) and the FP7/2007-2103 European Union project CVgenes@target (grant agreement number Health-F2-

2013-601456) (to M Dichgans).

*VISP*: The GWAS component of the VISP study was supported by the United States National Human Genome Research Institute (NHGRI), Grant U01 HG005160 (PI Michèle Sale & Bradford Worrall), as part of the Genomics and Randomized Trials Network (GARNET). Genotyping services were provided by the Johns Hopkins University Center for Inherited Disease Research (CIDR), which is fully funded through a federal contract from the NIH to the Johns Hopkins University. Assistance with data cleaning was provided by the GARNET Coordinating Center (U01 HG005157; PI Bruce S Weir). Study recruitment and collection of datasets for the VISP clinical trial were supported by an investigator-

initiated research grant (R01 NS34447; PI James Toole) from the United States Public Health Service, NINDS, Bethesda, Maryland. Control data for comparison with European ancestry VISP stroke cases were obtained through the database of genotypes and phenotypes (dbGAP) High Density SNP Association Analysis of Melanoma: Case-Control and Outcomes Investigation (phs000187.v1.p1; R01CA100264, 3P50CA093459, 5P50CA097007, 5R01ES011740, 5R01CA133996, HHSN268200782096C; PIs Christopher Amos, Qingyi Wei, Jeffrey E. Lee). For VISP stroke cases of African ancestry, a subset of the Healthy Aging in Neighborhoods of Diversity across the Life Span study (HANDLS) were used as stroke free controls. HANDLS is funded by the National Institute of Aging (1Z01AG000513; PI Michele K. Evans).

*WHI*: Funding support for WHI-GARNET was provided through the NHGRI GARNET (Grant Number U01 HG005152). Assistance with phenotype harmonisation and genotype cleaning, as well as with general study coordination, was provided by the GARNET Coordinating Center (U01 HG005157). Funding support for genotyping, which was performed at the Broad Institute of MIT and Harvard, was provided by the NIH Genes, Environment, and Health Initiative (GEI; U01 HG004424).

The SiGN study was funded by a cooperative agreement grant from the US National Institute of Neurological Disorders and Stroke, National Institutes of Health (U01 NS069208). The information about funding for each collection is reported below.

#### Discovery Case-only & Case and Control Cohorts:

**BASICMAR:** The Base de Datos de Ictus del Hospital del Mar (BASICMAR) Genetic Study was supported by the Ministerio de Sanidad y Consumo de España, Instituto de Salud Carlos III (ISC III) with the grants: Registro BASICMAR Funding for Research in Health (PI051737); GWA Study of Leukoaraiosis (GWALA) project from Fondos de Investigación Sanitaria ISC III (PI10/02064) and (PI12/01238); Agència de Gestió Ajuts Universitaris de Recerca (2014 SGR 1213) and Fondos European Regional Development Funding (FEDER/EDRF) Red INVICTUS-PLUS (RD16/0019/0002). Additional support was provided by the Fundació la Marató TV3 with the grant GODS project. Genestroke Consortium (76/C/2011) Recercaixa'13 (JJ086116). Assistance with data cleaning was provided by the Research in Cardiovascular and Inflammatory Diseases Program of Institute Hospital del Mar of Medical Investigations, Hospital del Mar, and the Barcelona Biomedical Research Park.

**EDINBURGH:** The Edinburgh Stroke Study was supported by the Wellcome Trust and the Binks Trust. Sample processing occurred in the Genetics Core Laboratory of the Wellcome Trust Clinical Research Facility, Western General Hospital, Edinburgh, UK. Much of the neuroimaging occurred in the Scottish Funding Council Brain Imaging Research Centre ([www.sbirc.ed.ac.uk](http://www.sbirc.ed.ac.uk)), University of Edinburgh, a core area of the Wellcome Trust Clinical Research Facility and part of the Scottish Imaging Network–A Platform for Scientific Excellence (SINAPSE) collaboration ([www.sinapse.ac.uk](http://www.sinapse.ac.uk)), funded by the Scottish Funding Council and the Chief Scientist Office. Genotyping was performed at the Wellcome Trust Sanger Institute in the United Kingdom and funded by the Wellcome Trust as part of the Wellcome Trust Case Control Consortium 2 project (085475/B/08/Z and 085475/Z/08/Z and WT084724MA).

**GCKNSS:** The Greater Cincinnati/Northern Kentucky Stroke Study (GCKNSS) was supported by the NIH (NS030678).

**GRAZ:** The Austrian Stroke Prevention Study was supported by the Austrian Science Fund (FWF) grant Nos. P20545-P05 and P13180 and I904-B13 (Era-Net). The Medical University of Graz supports the databases of the Graz Stroke Study and the Austrian Stroke Prevention Study.

**KRAKOW:** Phenotypic data and genetic specimens collection were funded by the grant from the Polish Ministry of Science and Higher Education for Leading National Research Centers (KNOW) and by the grants from the Jagiellonian University Medical College in Krakow, Poland: K/ZDS/002848, K/ZDS/003844.

**LEUVEN:** The Leuven Stroke genetics study was supported by personal research funds from the Department of Neurology of the University Hospitals Leuven. Dr Lemmens is a Senior Clinical Investigator of FWO Flanders (FWO 1841913N).

**LUND:** The Lund Stroke Register was supported by the Swedish Research Council (K2010-61X-20378-04-3), The Swedish Heart-Lung Foundation, Region Skåne, Skåne University Hospital, the Freemasons Lodge of Instruction EOS in Lund, King Gustaf V's and Queen Victoria's Foundation, Lund University, and the Swedish Stroke Association. Biobank services were provided by Region Skåne Competence Centre (RSKC Malmö), Skåne University Hospital, Malmö, Sweden, and Biobank, Labmedicin Skåne, University and Regional Laboratories Region Skåne, Sweden.

MALMÖ: The Malmö Diet and Cancer Study was supported by the Swedish Research Council (Vetenskapsrådet), Heart and Lung Foundation (Hjärt och Lungfonden), and Swedish Stroke Foundation (Strokeförbundet).

MCISS: The Middlesex County Ischemic Stroke Study (MCISS) was supported by intramural funding from the New Jersey Neuroscience Institute/JFK Medical Center, Edison, NJ, and The Neurogenetics Foundation, Cranbury, NJ. We acknowledge Dr Souvik Sen for his advice and encouragement in the initiation and design of this study.

MIAMISR and NOMAS: The Northern Manhattan Study (NOMAS) was supported by grants from the NINDS (R37 NS029993, R01 NS27517). The Cerebrovascular Biorepository at University of Miami/Jackson Memorial Hospital (The Miami Stroke Registry, Institutional Review Board No.

20070386) was supported by the Department of Neurology at University of Miami Miller School of

Medicine and Evelyn McKnight Brain Institute. Biorepository and DNA extraction services were provided by the Hussmann Institute for Human Genomics at the Miller School of Medicine.

MGH-GASROS: The Massachusetts General Hospital Stroke Genetics Group was supported by the NIH Genes Affecting Stroke Risks and Outcomes Study (GASROS) grant K23 NS042720, the American Heart Association/Bugher Foundation Centers for Stroke Prevention Research 0775010N, and NINDS K23NS042695, K23 NS064052, the Deane Institute for Integrative Research in Atrial Fibrillation and Stroke, and by the Keane Stroke Genetics Fund. Genotyping services were provided by the Broad Institute Center for Genotyping and Analysis, supported by grant U54 RR020278 from the National Center for Research Resources.

NHS: The Nurses' Health Study work on stroke is supported by grants from the NIH, including HL088521 and HL34594 from the National Heart, Lung, and Blood Institute, as well as grants from the National Cancer Institute funding the questionnaire follow-up and blood collection: CA87969 and CA49449.

OXVASC: The Oxford Vascular Study was supported by the Wellcome Trust, Wolfson Foundation, Stroke Association, Medical Research Council, Dunhill Medical Trust, NIH Research (NIHR), and NIHR Oxford Biomedical Research Centre based at Oxford University Hospitals NHS Trust and University of Oxford. Dr Rothwell is in receipt of Senior Investigator Awards from the Wellcome Trust and the NIHR.

REGARDS: The Reasons for Geographic and Racial Differences in Stroke (REGARDS) Study was supported by a cooperative agreement U01 NS041588 from the NINDS, NIH, and Department of Health and Human Service. A full list of participating REGARDS investigators and institutions can be found at <http://www.regardsstudy.org>.

SPS3: The Secondary Prevention of Small Subcortical Strokes trial was funded by the US National Institute of Health and Neurological Disorders and Stroke grant No. U01NS38529-04A1 (principal investigator, Oscar R. Benavente; coprincipal investigator, Robert G. Hart). The SPS3 Genetic Substudy (SPS3-GENES) was funded by R01 NS073346 (coprincipal investigators, Julie A. Johnson, Oscar R. Benavente, and Alan R. Shuldiner) and U01 GM074492-05S109 (principal investigator, Julie A. Johnson).

ST. GEORGE'S: The principal funding for this study was provided by the Wellcome Trust, as part of the Wellcome Trust Case Control Consortium 2 project (085475/B/08/Z and 085475/Z/08/Z and WT084724MA). Collection of some of the St George's stroke cohort was supported by project grant support from the Stroke Association. Hugh Markus is supported by an NIHR Investigator award. Matthew Traylor was supported by project grant funding from the Stroke Association.(TSA 2013/01)

VISP: The GWAS component of the VISP study was supported by the United States National Human Genome Research Institute (NHGRI), Grant U01 HG005160 (PI Michèle Sale & Bradford Worrall), as part of the Genomics and Randomized Trials Network (GARNET). Genotyping services were provided by the Johns Hopkins University Center for Inherited Disease Research (CIDR), which is fully funded through a federal contract from the NIH to the Johns Hopkins University. Assistance with data cleaning was provided by the GARNET Coordinating Center (U01 HG005157; PI Bruce S Weir). Study recruitment and collection of datasets for the VISP clinical trial were supported by an investigator- initiated research grant (R01 NS34447; PI James Toole) from the United States Public Health Service, NINDS, Bethesda, Maryland. Control data for comparison with European ancestry VISP stroke cases were obtained through the database of genotypes and phenotypes (dbGAP) High Density SNP Association Analysis of Melanoma: Case-Control and Outcomes Investigation (phs000187.v1.p1; R01CA100264, 3P50CA093459, 5P50CA097007, 5R01ES011740, 5R01CA133996, HHSN268200782096C; PIs Christopher Amos, Qingyi Wei, Jeffrey E. Lee). For VISP stroke cases of African ancestry, a subset of the Healthy Aging in Neighborhoods of Diversity across the Life Span study (HANDLS) were used as stroke free controls. HANDLS is funded by the National Institute of Aging (1Z01AG000513; PI Michele K. Evans).

WHI-OS: The Women's Health Initiatives (WHI) program was funded by the National Heart, Lung, and Blood Institute, NIH, US Department of Health and Human Services through contracts N01WH22110, 24152, 32100-2, 32105-6, 32108-9, 32111-13, 32115, 32118 to 32119, 32122, 42107-26, 42129-32, and 44221. The Hormones and Biomarkers Predicting Stroke (HaBPS) was supported by a grant from the National Institutes of Neurological Disorders and Stroke (R01NS042618).

WUSTL: Washington University St. Louis Stroke Study (WUSTL): The collection, extraction of DNA from blood, and storage of specimens were supported by 2 NINDS NIH grants (P50 NS055977 and R01 NS8541901). Basic demographic and clinical characterization of stroke phenotype was prospectively collected in the Cognitive Rehabilitation and Recovery Group (CRRG) registry. The Recovery Genomics after Ischemic Stroke (ReGenesIS) study was supported by a grant from the Barnes-Jewish Hospital Foundation.

Control-only  
Cohorts:

ADHD: The ADHD-VHIR study from Barcelona was supported by Fundació Marató de TV3 (ref. 092330/31), the Miguel Servet Programme, Instituto de Salud Carlos III (CP09/00119 and CPII15/00023) and the European Regional Development Fund (ERDF).

Health ABC: The Health Aging and Body Composition Study was supported by NIA contracts N01AG62101, N01AG62103, and N01AG62106 and, in part, by the NIA Intramural Research Program. The genome-wide association study was funded by NIA grant 1R01AG032098-01A1 to Wake Forest University Health Sciences and genotyping services were provided by the Center for Inherited Disease Research (CIDR). CIDR is fully funded through a federal contract from the National Institutes of Health to The Johns Hopkins University, contract number HHSN268200782096C. This study utilized the high-performance computational capabilities of the Biowulf Linux cluster at the National Institutes of Health, Bethesda, Md. (<http://biowulf.nih.gov>). Mike A. Nalls' participation is supported by a consulting contract between Data Tecnica International (Dr. Nalls is the founder and primary consultant) and the National Institute on Aging, NIH, Bethesda, MD, USA, as a possible conflict of interest Dr. Nalls also consults for Illumina Inc, the Michael J. Fox Foundation and University of California Healthcare.

HCHS/SOL: The Hispanic Community Health Study/Study of Latinos was carried out as a collaborative study supported by contracts from the National Heart, Lung, and Blood Institute (NHLBI) to the University of North Carolina (N01-HC65233), University of Miami (N01-HC65234), Albert Einstein College of Medicine (N01-HC65235), Northwestern University (N01-HC65236), San Diego State University (N01-HC65237), and University of Washington (HHSN268201300005C). The following Institutes/Centers/Offices contribute to the HCHS/SOL through a transfer of funds to the NHLBI: National Center on Minority Health and Health Disparities, the National Institute of Deafness and Other Communications Disorders, the National Institute of Dental and Craniofacial Research, the National Institute of Diabetes and Digestive and Kidney Diseases, the National Institute of Neurological Disorders and Stroke, and the Office of Dietary Supplements.

The authors thank the staff and participants of HCHS/SOL for their important contributions. A complete list of staff and investigators has been provided by Sorlie P., et al. in *Ann Epidemiol.* 2010 Aug;20: 642-649 and is also available on the study website <http://www.csc.unc.edu/hchs/>.

HRS: HRS is supported by the National Institute on Aging (NIA U01AG009740). The genotyping was funded as a separate award from the National Institute on Aging (RC2 AG036495). Genotyping was conducted by the NIH Center for Inherited Disease Research (CIDR) at Johns Hopkins University. Genotyping quality control and final preparation of the data were performed by the Genetics Coordinating Center at the University of Washington. HRS genotype data have been deposited in the NIH GWAS repository (dbGaP). Researchers wishing to use the HRS genetic data must first apply to dbGaP for access. The process to request access to any dbGaP study is done via the dbGaP authorized access system. Researchers who wish to obtain HRS phenotype measures that are not in dbGaP must submit a data access use agreement to HRS. For further information, contact [hrsquestions@umich.edu](mailto:hrsquestions@umich.edu).

Relevant websites describing HRS genotype and phenotype data are: [www.ncbi.nlm.nih.gov/projects/gap/cgi-](http://www.ncbi.nlm.nih.gov/projects/gap/cgi-) and <http://hrsonline.isr.umich.edu>.

INMA: This study was funded by grants from Instituto de Salud Carlos III (CB06/02/0041, G03/176, FIS PI041436, PI081151, PI041705, PI061756, PI091958, and PS09/00432, FIS-FEDER 03/1615, 04/1509, 04/1112, 04/1931, 05/1079, 05/1052, 06/1213, 07/0314, 09/02647, 11/01007, 11/02591, 11/02038, 13/1944, 13/2032 and CP11/0178), Spanish Ministry of Science and Innovation (SAF2008-00357), European Commission (ENGAGE project and grant agreement HEALTH-F4-2007-201413, HEALTH.2010.2.4.5-1, FP7-ENV-2011 cod 282957), Fundació La Marató de TV3, Generalitat de Catalunya-CIRIT 1999SGR 00241 and Conselleria de Sanitat Generalitat Valenciana. Part of the DNA extractions and genotyping was performed at the Spanish National Genotyping Centre (CEGEN- Barcelona). The authors are grateful to Silvia Fochs, Anna Sánchez, Maribel López, Nuria Pey, Muriel Ferrer, Amparo Quiles, Sandra Pérez, Gemma León, Elena Romero, Maria Andreu, Nati Galiana, Maria Dolores Climent, Amparo Cases and Cristina Capo for their assistance in contacting the families and administering the questionnaires. The authors would particularly like to thank all the participants for their generous collaboration. A full roster of the INMA Project Investigators can be found at [http://www.proyectoinma.org/presentacion-inma/listado-investigadores/en\\_listado-investigadores.html](http://www.proyectoinma.org/presentacion-inma/listado-investigadores/en_listado-investigadores.html).

KORA: The KORA research platform (KORA, Cooperative Research in the Region of Augsburg) was initiated and financed by the Helmholtz Zentrum München - German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research and by the State of Bavaria. Furthermore, KORA research was supported within the Munich Center of Health Sciences (MC Health), Ludwig-Maximilians-Universität, as part of LMUinnovativ.

OAI: The OAI is a public-private partnership comprised of five contracts (N01-AR-2-2258; N01-AR-2-2259; N01-AR-2-2260; N01-AR-2-2261; N01-AR-2-2262) funded by the National Institutes of Health, a branch of the Department of Health and Human Services, and conducted by the OAI Study Investigators. Genotyping support was provided by grant RC2-AR-058950 from NIAMS/NIH. Private funding partners include Merck Research Laboratories; Novartis Pharmaceuticals Corporation, GlaxoSmithKline; and Pfizer, Inc. Private sector funding for the OAI is managed by the Foundation for the National Institutes of Health.

WTCCC2: Wellcome Trust Case-Control Consortium 2 (WTCCC2) was principally funded by the Wellcome Trust, as part of the Wellcome Trust Case Control Consortium 2 project (085475/B/08/Z and 085475/Z/08/Z and WT084724MA). The Stroke Association provided additional support for collection of some of the St George's, London cases. The Oxford cases were collected as part of the Oxford Vascular Study which is funded by the MRC, Stroke Association, Dunhill Medical Trust, National

Institute of Health Research (NIHR) and the NIHR Biomedical Research Centre, Oxford. The Edinburgh Stroke Study was supported by the Wellcome Trust (clinician scientist award to C Sudlow), and the Binks Trust. Sample processing occurred in the Genetics Core Laboratory of the Wellcome Trust Clinical Research Facility, Western General Hospital, Edinburgh. Much of the neuroimaging occurred in the Scottish Funding Council Brain Imaging Research Centre ([www.sbirc.ed.ac.uk](http://www.sbirc.ed.ac.uk)), Division of Clinical Neurosciences, University of Edinburgh, a core area of the Wellcome Trust Clinical Research Facility and part of the SINAPSE (Scottish Imaging Network—A Platform for Scientific Excellence) collaboration ([www.sinapse.ac.uk](http://www.sinapse.ac.uk)), funded by the Scottish Funding Council and the Chief Scientist Office. Collection of the Munich cases and data analysis was supported by the Vascular Dementia Research Foundation. M Farrall and A Helgadottir acknowledge support from the BHF Centre of Research Excellence in Oxford and the Wellcome Trust core award (090532/Z/09/Z). Barcelona The Neurovascular Research Laboratory takes part in the International Stroke Genetics Consortium (ISGC), the Spanish Stroke Genetics Consortium ([www.genestroke.com](http://www.genestroke.com)), and the Cooperative Neurovascular Research RENEVAS (RD06/0026/0010). This study was funded by a grant of the Spanish government (PI10/01212.). The research leading to these results has received funding from the European Union's Seventh Framework Programme (FP7/2007-2013) under grant agreements

#201024 and #202213 (European Stroke Network). Belgium Stroke Study (BSS) was supported by Erasme Funds. Edinburgh Stroke Study (ESS) (which contributed discovery cases as part of WTCCC2 and additional replication cases) was supported as described above. Lothian Birth Cohort 1936 was supported in part by Research into Aging, Help the Aged (Sidney De Haan Award and The Disconnected Mind Major Gift Campaign), MRC, and UK Biotechnology and Biological Sciences Research Council (BBSRC). Lothian Birth Cohort 1936 was also supported by a programme grant from Research Into Ageing and continues with programme grants from Help the Aged/Research Into Ageing (Disconnected Mind). The work was undertaken by The University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology, part of the cross council Lifelong Health and Wellbeing Initiative (G0700704/84698). Funding from the BBSRC, Engineering and Physical Sciences Research Council (EPSRC), Economic and Social Research Council (ESRC), and MRC is gratefully acknowledged. Genotyping of the LBC1936 was funded by the BBSRC.

Secondary  
Cohorts:

INTERSTROKE: We would like to acknowledge the Canadian Institutes of Health Research, Heart and Stroke Foundation of Canada, Canadian Stroke Network, Pfizer Cardiovascular Award, Merck, AstraZeneca, and Boehringer Ingelheim.

MILANO: Milano- Besta Stroke Register Collection and genotyping of the Milan cases within CEDIR were supported by Annual Research Funding of the Italian Ministry of Health (Grant Numbers: RC 2007/LR6, RC 2008/LR6; RC 2009/LR8; RC 2010/LR8). FP6 LSHM-CT-2007-037273 for the PROCARDIS control samples.

PROSPER: The Prospective Study on Pravastatin in the Elderly at Risk (PROSPER) was supported by an investigator initiated grant obtained from Bristol-Myers Squibb. Prof. Dr. J. W. Jukema is an Established Clinical Investigator of the Netherlands Heart Foundation (grant 2001 D 032). Support for genotyping was provided by the seventh framework program of the European commission (grant 223004) and by the Netherlands Genomics Initiative (Netherlands Consortium for Healthy Aging grant 050-060-810).

RACE: We are thankful to the RACE study participants. Fieldwork in RACE was funded by the R-21 grant provided by the NINDS and the Fogarty International Center (1R21NS064908-01) and educational grants available to Dr. Saleheen at the Center for Non-Communicable Diseases, Pakistan. We would also like to acknowledge the contributions made by Professor John Danesh, Dr. Ayeesha Kamal and Professor Panos Deloukas.

SIGNET: The Sea Islands Genetics Network (SIGNET) was supported by R01 DK084350 (MM Sale) from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), and consists of data from the REasons for Geographic And Racial Differences in Stroke (REGARDS) cohort, (U01 NS041588; G Howard), Project SuGAR (Sea Islands Genetic African American Registry) (W.M. Keck Foundation; WT Garvey), a South Carolina Center of Biomedical Research Excellence (COBRE) in Oral Health Project P20 RR017696 (PI: Kirkwood; Sub-award: JK Fernandes), and the Systemic Lupus Erythematosus in Gullah Health (SLEIGH) study (PI: GS Gilkeson; K23 AR052364, DL Kamen; UL1 RR029882, KT Brady). Only data from the SIGNET-REGARDS sub-study were included in the current analyses. REGARDS is supported by a cooperative agreement U01 NS041588 from the National Institute of Neurological Disorders and Stroke (NINDS), National Institutes of Health (NIH), Department of Health and Human Service. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIDDK, NINDS or the NIH. Representatives of the funding agencies have been involved in the review of the manuscript but not directly involved in the collection, management, analysis or interpretation of the data. The authors thank the other investigators, the staff, and the participants of the REGARDS study for their valuable contributions. A full list of participating REGARDS investigators and institutions can be found at <http://www.regardsstudy.org>

Utrecht ImmunoChip/PROMISe: S.Achterberg was supported by in part by a grant from the Netherlands Heart Foundation, (grant no 2005B031) and a grant from the Dutch Brain Foundation (project 2008(1).10).



VHIR-FMT-Barcelona: The Barcelona GWAs Study was supported by the Genetic contribution to functional Outcome and Disability after Stroke (GODS) project, Fundació la Marató de TV3, GENERACION Project (Instituto de Salud Carlos III: PI15/01978), Pre-Test Stroke Project (Instituto de Salud Carlos III: PMP15/00022) and by the Miguel Servet grant (Pharmastroke project: CP12/03298). Neurovascular research Laboratory and the Stroke Pharmacogenomics and Genetics lab take part in the INVICTUS+ network. I. F-C. is supported by the Miguel Servet programme (CP12/03298), Instituto de Salud Carlos III.

WGHS: The WGHS is supported by the National Heart, Lung, and Blood Institute (HL043851, HL080467, and HL099355) and the National Cancer Institute (CA047988 and UM1CA182913), with collaborative scientific support and funding for genotyping provided by Amgen.

WHI-HT: WHI Funding support for WHI-GARNET was provided through the NHGRI GARNET (Grant Number U01 HG005152). Assistance with phenotype harmonisation and genotype cleaning, as well as with general study coordination, was provided by the GARNET Coordinating Center (U01 HG005157). Funding support for genotyping, which was performed at the Broad Institute of MIT and Harvard, was provided by the NIH Genes, Environment, and Health Initiative (GEI; U01 HG004424).

## **CHARGE**

*AGES*: Age, Gene/Environment Susceptibility (AGES) -Reykjavik The study was funded by the National Institute on Aging (NIA)(N01-AG-12100), Hjartavernd (the Icelandic Heart Association), and the Althingi (the Icelandic Parliament), with contributions from the Intramural Research Programs at the NIA, the National Heart, Lung, and Blood Institute, and the National Institute of Neurological Disorders and Stroke (Z01 HL004607-08 CE).

*ARIC*: The Atherosclerosis Risk in Communities study was performed as a collaborative study supported by National Heart, Lung, and Blood Institute (NHLBI) contracts (HHSN268201100005C, HSN268201100006C, HSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C), R01HL70825, R01HL087641, R01HL59367, and R01HL086694; National Human Genome Research Institute contract U01HG004402; and National Institutes of Health (NIH) contract HHSN268200625226C. Infrastructure was partly supported by grant No. UL1RR025005, a component of the NIH and NIH Roadmap for Medical Research. This project was also supported by NIH R01 grant NS087541 to MF. *CHS*: This CHS research was supported by NHLBI contracts HHSN268201200036C, HHSN268200800007C, N01HC55222, N01HC85079, N01HC85080, N01HC85081, N01HC85082, N01HC85083, N01HC85086, and HHSN268200960009C; and NHLBI grants U01HL080295, R01HL087652, R01HL105756, R01HL103612, R01HL120393, and R01HL130114 with additional contribution from the National Institute of Neurological Disorders and Stroke (NINDS). Additional support was provided through R01AG023629 from the National Institute on Aging (NIA). A full list of principal CHS investigators and institutions can be found at CHS-NHLBI.org. The provision of genotyping data was supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR000124, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

*FHS*: This work was supported by the National Heart, Lung and Blood Institute's Framingham Heart Study (Contract No. N01-HC-25195 and No. HHSN268201500001I) and its contract with Affymetrix, Inc. for genotyping services (Contract No. N02-HL-6-4278). A portion of this research utilized the Linux Cluster for Genetic Analysis (LinGA-II) funded by the Robert Dawson Evans Endowment of the Department of Medicine at Boston University School of Medicine and

Boston Medical Center. This study was also supported by grants from the National Institute of Neurological Disorders and Stroke (R01 NS017950), the National Heart, Lung and Blood Institute (R01 HL093029) and the National Institute of Aging (R01s AG033193, AG008122, AG016495, U01 AG049505).

*FINRISK*: V.S. was supported by the Academy of Finland (grant #139635) and the Finnish Foundation for Cardiovascular Research

*HEALTH ABC*: The Health ABC Study was supported by NIA contracts N01AG62101, N01AG62103, and N01AG62106 and, in part, by the NIA Intramural Research Program. The genome-wide association study was funded by NIA grant 1R01AG032098-01A1 to Wake Forest University Health Sciences and genotyping services were provided by the Center for Inherited Disease Research (CIDR). CIDR is fully funded through a federal contract from the National Institutes of Health to The Johns Hopkins University, contract number HHSN268200782096C. This study utilized the high-performance computational capabilities of the Biowulf Linux cluster at the National Institutes of Health, Bethesda, Md. (<http://biowulf.nih.gov>).

*Rotterdam Study*: The generation and management of GWAS genotype data for the Rotterdam Study is supported by the Netherlands Organisation of Scientific Research NWO Investments (nr. 175.010.2005.011, 911-03-012). This study is funded by the Research Institute for Diseases in the Elderly (014-93-015; RIDE2), the Netherlands Genomics Initiative (NGI)/Netherlands Organisation for Scientific Research (NWO) project nr. 050-060-810. The Rotterdam Study is funded by Erasmus

Medical Center and Erasmus University, Rotterdam, Netherlands Organization for the Health Research and Development (ZonMw), the Research Institute for Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam. MAI is supported by an NWO Veni grant (916.13.054). *SHIP*: SHIP is part of the Community Medicine Research net of the University of Greifswald, Germany, which is funded by the Federal State of Mecklenburg-West Pomerania. The data collections underlying this work have been funded by the Federal Ministry of Education and Research (grants no. 01ZZ9603, 01ZZ0103, and 01ZZ0403), the Ministry of Cultural Affairs as well as the Social Ministry of the Federal State of Mecklenburg-West Pomerania, and the German Research Foundation (SCHM 2744/1-1). Genome-wide data have been supported by the Federal Ministry of Education and Research (grant no. 03ZIK012) and a joint grant from Siemens Healthcare, Erlangen, Germany and the Federal State of Mecklenburg- West Pomerania.

*WGHS*: The WGHS is supported by HL043851, HL080467, and HL099355. from the National Heart, Lung, and Blood Institute and CA047988 from the National Cancer Institute with collaborative scientific support and funding for genotyping provided by Amgen.

*MESA*: MESA and the MESA SHARe project are conducted and supported by the National Heart,

Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts HHSN268201500003I, N01-HC-95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-001079, UL1-TR-001420, UL1-TR-001881, and DK063491. Funding for SHARe genotyping was provided by NHLBI Contract N02-HL-64278. Genotyping was performed at Affymetrix (Santa Clara, California, USA) and the Broad Institute of Harvard and MIT (Boston, Massachusetts, USA) using the Affymetrix Genome-Wide Human SNP Array 6.0.

*PROSPER*: The PROSPER study was supported by an investigator initiated grant obtained from Bristol-Myers Squibb. Prof. Dr. J. W. Jukema is an Established Clinical Investigator of the Netherlands Heart Foundation (grant 2001 D 032). Support for genotyping was provided by the seventh framework program of the European commission (grant 223004) and by the

Netherlands Genomics Initiative (Netherlands Consortium for Healthy Aging grant 050-060-810).

**TWINGENE:** This work was supported by grants from the Ministry for Higher Education, the Swedish

Research Council (M-2005-1112 and 2009-2298), GenomeUtwinn (EU/QLRT-2001-01254; QLG2-CT-

2002-01254), NIH grant DK U01-066134, The Swedish Foundation for Strategic Research (SSF; ICA08-0047).

**ULSAM:** This work was supported by grants from Uppsala University, Swedish Research Council (2012-1397), Swedish Heart-Lung Foundation (20140422), Knut och Alice Wallenberg Foundation, European Research Council, Swedish Diabetes Foundation (2013-024). Andrew P Morris is a Wellcome Trust Senior Fellow in Basic Biomedical Science (grant numbers WT064890, WT090532 and WT098017)

**3C:** The 3-City Study is conducted under a partnership agreement among the Institut National de la Santé et de la Recherche Médicale (INSERM), the University of Bordeaux, and Sanofi-Aventis. The Fondation pour la Recherche Médicale funded the preparation and initiation of the study. The 3C Study is also supported by the Caisse Nationale Maladie des Travailleurs Salariés, Direction Générale de la Santé, Mutuelle Générale de l'Éducation Nationale (MGEN), Institut de la Longévité, Conseils Régionaux of Aquitaine and Bourgogne, Fondation de France, and Ministry of Research-INSERM Programme "Cohortes et collections de données biologiques." This work was supported by the National Foundation for Alzheimer's Disease and Related Disorders, the Institut Pasteur de Lille, the Centre National de Génotypage and the LABEX (Laboratory of Excellence program investment for the future) DISTALZ - Development of Innovative Strategies for a Transdisciplinary approach to Alzheimer's

disease. Stéphanie Debette is supported by a grant from the Fondation Leducq, a starting grant from the European Research Council, a grant from the Joint Programme of Neurodegenerative Disease research, and the Initiative of Excellence of Bordeaux University.

**EPIC:** EPIC was funded by the UK Medical Research Council (G0800270), British Heart Foundation (SP/09/002), UK National Institute for Health Research Cambridge Biomedical Research Centre, European Research Council (268834), European Commission Framework Programme 7 (HEALTH- F2-2012-279233).

**AIDHS/SDS:** This work was supported by NIH grants -R01DK082766 funded by the National Institute of Health (NIDDK) and NOT-HG-11-009 funded by NHGRI, VPR Bridge Grant and Harold Hamm Enrichment Grants from University of Oklahoma Health Sciences Center. Authors thank all the participants of AIDHS/SDS and are grateful for their contribution in this study.

**VHIR-FMT-Barcelona:** The Barcelona GWAs Study was supported by the Genetic contribution to functional Outcome and Disability after Stroke (GODS) project, Fundació la Marató de TV3 and by the Miguel Servet grant (Pharmastroke project: CP12/03298). Neurovascular research Laboratory takes part in the INVICTUS network. I. F-C. is supported by the Miguel Servet programme (CP12/03298), Instituto de Salud Carlos III.

#### **Biobank Japan:**

We would like to express our greatfulness to all the members of J-MICC, JPHC, and TMM. We extend our appreciation to staffs of BBJ for their outstanding assistance. This study was funded by the BioBank Japan project, which is supported by the Ministry of Education, Culture, Sports, Sciences and Technology (MEXT) of Japanese government and the Japan Agency for Medical Research and Development (AMED).

**CADISP:** The Cervical Artery Dissections and Ischemic Stroke Patients (CADISP) study has been supported by Inserm, Lille 2 University, Institut Pasteur de Lille and Lille University Hospital and received funding from the ERDF (FEDER funds) and Région Nord-Pas de Calais in the

frame of Contrat de Projets Etat-Region 2007-2013 Région Nord-Pas-de-Calais - Grant N°09120030, Centre National de Genotypage, Emil Aaltonen Foundation, Paavo Ilmari Ahvenainen Foundation, Helsinki University Central Hospital Research Fund, Helsinki University Medical Foundation, Päivikki and Sakari Sohlberg Foundation, Aarne Koskelo Foundation, Maire Taponen Foundation, Aarne and Aili Turunen Foundation, Lilly Foundation, Alfred Kordelin Foundation, Finnish Medical Foundation, Orion Farnos Research Foundation, Maud Kuistila Foundation, the Finnish Brain Foundation, Biomedicum Helsinki Foundation, Projet Hospitalier de Recherche Clinique Régional, Fondation de France, Génopôle de Lille, Adrinord, Basel Stroke-Funds, Käthe-Zingg-Schwichtenberg-Fonds of the Swiss Academy of Medical Sciences, Swiss Heart Foundation.

### **COMPASS:**

*CHS:* The CHS study was supported by National Heart, Lung, and Blood Institute contracts HHSN268201200036C, HHSN268200800007C, N01HC55222, N01HC85079, N01HC85080, N01HC85081, N01HC85082, N01HC85083, N01HC85086; and NHLBI grants U01HL080295, R01HL087652, R01HL105756, R01HL103612, R01HL120393, R01HL130114, and R01HL085251 with contributions from the National Institute of Neurological Disorders and Stroke. Additional support

was provided through R01AG023629 from the National Institute on Aging. A full list of principal CHS investigators and institutions can be found at CHS-NHLBI.org. The provision of genotyping data were supported, in part, by the National Center for Advancing Translational Sciences, grant UL1TR000124, and the National Institute of Diabetes and Digestive and Kidney Disease Research Center grant DK063491 to the Southern California Diabetes Endocrinology Research Center. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

*HANDLS:* Healthy Aging in Neighborhoods of Diversity across the Life Span was supported by the Intramural Research Program of the NIH, National Institute of Aging and the National Center on Minority Health and Health Disparities (project no. Z01-AG000513 and human subjects protocol no.

2009-149).

*INTERSTROKE:* The INTERSTROKE study was supported by the Canadian Institutes of Health Research, Heart and Stroke Foundation of Canada, Canadian Stroke Network, Health Research Board Ireland, Swedish Research Council, Swedish Heart and Lung Foundation, The Health & Medical Care Committee of the Regional Executive Board, Region Västra Götaland (Sweden), AstraZeneca, Boehringer Ingelheim (Canada), Pfizer (Canada), MSD, Chest, Heart and Stroke Scotland, and The Stroke Association, with support from The UK Stroke Research Network. Microarray genotyping for a subset of INTERSTROKE participants was funded by the Heart and Stroke Foundation of Canada Grant NA-6872 (PI: Drs Pare, Anand, O'Donnell, Xie, Yusuf).

*ISGS and SWISS:* Ischemic Stroke Genetics Study (ISGS) and Siblings with Ischemic Stroke Study (SWISS) were supported by the National Institute of Neurological Disorders and Stroke grants (R01 NS42733; PI Meschia) and (R01NS39987; PI Meschia), respectively with additional support, in part, from the Intramural Research Program of the National Institute of Aging (Z01 AG000954-06; PI Singleton). Both studies used samples and clinical data from the NIH-NINDS Human Genetics Resource Center DNA and Cell Line Repository, human subjects protocol no. 2003-081 and 2004-147.

**SIGNET-REGARDS:** We would like to thank all of the participants of the REGARDS Study for their valuable contributions, as well as REGARDS investigators and staff. The Sea Islands Genetics Network (SIGNET) was supported by R01 DK084350 (MM Sale), and SIGNET-REGARDS consists of data from the REasons for Geographic And Racial Differences in Stroke (REGARDS) cohort, supported by a cooperative agreement U01 NS041588 (G Howard).

**VISP:** Vitamin Intervention for Stroke Prevention (VISP) was funded by the National Institute of Neurological Disorders and Stroke (R01-NS34447). Genome-wide association study data for a subset of VISP participants supported by the National Human Genome Research Institute (U01-HG005160), as part of the Genomics and Randomized Trials Network (PI: Drs Sale and Worrall).

**JHS:** The Jackson Heart Study is supported by contracts HHSN268201300046C, HHSN268201300047C, HHSN268201300048C, HHSN268201300049C, HSN268201300050C from the National Heart, Lung, and Blood Institute and the National Institute on Minority Health and Health Disparities. The authors also wish to thank the staffs and participants of the JHS. A full list of the participating institution and investigators can be found at <https://www.jacksonheartstudy.org/jhsinfo/Home/tabid/36/Default.aspx>.

**WHI:** The WHI program is funded by the National Heart, Lung, and Blood Institute, National Institutes of Health, U.S. Department of Health and Human Services through contracts HHSN268201600018C, HHSN268201600001C, HHSN268201600002C, HHSN268201600003C, and HHSN268201600004C.

The authors thank the WHI investigators and staff for their dedication and the study participants for making the program possible. A full listing of WHI investigators can be found at:

<http://www.whi.org/researchers/Documents%20%20Write%20a%20Paper/WHI%20Investigator%20Long%20List.pdf>.

**Glasgow Stroke Sample:** The work was supported by NHS Greater Glasgow Endowment funds.

**Helsinki 2000 Ischemic Stroke Genetics Study:** The study was supported by the Finnish Medical Foundation and the Helsinki University Central Hospital governmental subsidiary funds for clinical research. The investigators would like to thank Marja Metso, RN for her support of the study.

**Hisayama-ESR study:**

We thank Prof. Takanari Kitazono and Prof. Masahiro Kamouchi (Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan) for collecting clinical samples.

**HVH 1 & 2:**

The Heart and Vascular Health Study was supported by NHLBI grants R01HL085251, R01HL073410, and R01HL068986.

**INTERSTROKE:** We would like to acknowledge the Canadian Institutes of Health Research, Heart and Stroke Foundation of Canada, Canadian Stroke Network, Pfizer Cardiovascular Award, Merck, AstraZeneca, and Boehringer Ingelheim.

**MDC:** The Malmö Diet and Cancer Study was supported by the Swedish Research Council (Vetenskapsrådet), Heart and Lung Foundation (Hjärt och Lungfonden), and Swedish Stroke Foundation (Strokeförbundet).

**RACE:** Fieldwork in RACE was funded by the R-21 grant provided by the NINDS and the Fogarty International Center (1R21NS064908-01) and educational grants available to Dr.

Saleheen at the Center for Non-Communicable Diseases, Pakistan. We would also like to acknowledge the contributions made by Professor John Danesh, Dr. Ayeesha Kamal and Professor Panos Deloukas.

**SAHLSIS:** The Sahlgrenska Academy Study of Ischemic Stroke was supported by the Swedish Research Council (K2014-64X-14605-12-5), the Swedish Heart and Lung Foundation, the Swedish state/Sahlgrenska University Hospital (ALFGBG-429981), the Swedish Stroke Association, the Swedish Society of Medicine, and the Rune and Ulla Amlöv Foundation.

**SIFAP:** The sifap study (Stroke In Young Fabry Patients, <http://www.sifap.eu>; ClinicalTrials.gov: NCT00414583) has been supported partially by an unrestricted scientific grant from Shire Human Genetic Therapies. Funding for genotyping and analysis of samples were supported by the National Institutes of Health Genes, Environment and Health Initiative (GEI) Grant U01 HG004436, as part of the GENEVA consortium.

**SLESS:** This work was supported by a Stroke Association (UK) Programme Grant (PROG 3), the National Institute for Health Research Biomedical Research Centre (NIHR BRC) at South London and Maudsley NHS Foundation Trust and the National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London. Dr Rutten-Jacobs is supported by a British Heart Foundation Fellowship (FS/15/61/31626). Dr Markus is

supported by the National Institute for Health Research Cambridge University Hospitals Comprehensive Biomedical Research Centre and a National Institute for Health Research Senior Investigator award.

**UK - young lacunar stroke DNA resource:** Collection of the UK Young Lacunar Stroke DNA Study (DNA Lacunar) was primarily supported by the Wellcome Trust (WT072952) with additional support from the Stroke Association (TSA 2010/01). Genotyping of the samples, and Dr Traylor, were supported by a Stroke Association Grant (TSA 2013/01). Dr Markus is supported by the National Institute for Health Research Cambridge University Hospitals Comprehensive Biomedical Research Centre and a National Institute for Health Research Senior Investigator award. Matthew Traylor is funded by a BHF programme grant RG/16/4/32218. Loes Rutten-Jacobs is supported by a BHF Early career fellowship BHF/15/61/31626.

## **ICH**

Funding provided as follows: GERFHS, NIH grants NS36695 and NS30678; GOCHA, NIH grant R01NS059727, the Keane Stroke Genetics Research Fund, the Edward and Maybeth Sonn Research Fund, and the University of Michigan General Clinical Research Center M01 RR000042; ERICH, NIH grant NS069763; HM-ICH, Instituto de Salud Carlos III with the grants "Registro BASICMAR" Funding for Research in Health (PI051737), "GWALA project" from Fondos de Investigación Sanitaria ISC III (PI10/02064), and Fondos FEDER/EDRF Red de Investigación Cardiovascular (RD12/0042); JUHSS, Polish Ministry of Education grant N402 083934; LSR, Lund University, Region Skåne, the Swedish Research Council (K2010-61X-20378-04-3), the Swedish Stroke Association, the Freemasons Lodge of Instruction EOS in Lund, and the King Gustaf V and Queen Victoria's foundations; G.J.F. and H.B.B., NIH SPOTRIAS fellowship P50NS061343; C.D.A., fellowship from the American Brain Foundation; J.N.G., NIH grant 5K23NS059774; P.M.R., awards from the NIHR and the Wellcome Trust; M.S., NIH grant U01 NS074425; and D.L.B., NIH grants R01 NS062675, R01 HL098065, R01 NS070941, and R18 HS017690, the Blue Cross Blue Shield of Michigan Foundation, Michigan Department of Community Health, and the University of Michigan.

DAT and VC were financially supported by the Transatlantic Networks of Excellence Award (12CVD02) from Foundation Leducq

Statistical analyses were performed using the C2BIG computing cluster, funded by the Région Ile de France, Pierre and Marie Curie University, and the Institute for Cardiometabolism and Nutrition (ANR-10-IAHU-05).